



## Tansley review

# Grass secondary cell walls, *Brachypodium distachyon* as a model for discovery

Authors for correspondence:

Samuel P. Hazen

Tel: +1 413 545 4546

Email: hazensam@umass.edu

Richard Sibout

Tel: +33 2 40 67 50 65

Email: richard.sibout@inrae.fr

Received: 4 December 2019

Accepted: 5 March 2020

Joshua H. Coomey<sup>1,2</sup> , Richard Sibout<sup>3\*</sup>  and Samuel P. Hazen<sup>1,2\*</sup> 

<sup>1</sup>Biology Department, University of Massachusetts, Amherst, MA 01003, USA; <sup>2</sup>Plant Biology Graduate Program, University of Massachusetts, Amherst, MA 01003, USA; <sup>3</sup>Biopolymères Interactions Assemblages, INRAE, UR BIA, F-44316, Nantes, France

## Contents

Summary	1649	VIII. Hydroxycinnamic acids	1657
I. Introduction to the secondary cell wall	1650	IX. Tricin	1658
II. <i>Brachypodium distachyon</i> , a model grass system	1651	X. Silicon	1658
III. Cellulose	1652	XI. Transcriptional regulation of secondary cell wall thickening	1659
IV. Mixed-linkage glucans	1652	XII. Conclusions	1662
V. Grass heteroxylans	1654	Acknowledgements	1662
VI. Xylan acetylation	1655	References	1662
VII. Lignins	1655		

## Summary

*New Phytologist* (2020) **227**: 1649–1667  
doi: 10.1111/nph.16603

**Key words:** *Brachypodium distachyon*, grass secondary cell wall, hemicellulose, lignin, transcriptional regulation.

A key aspect of plant growth is the synthesis and deposition of cell walls. In specific tissues and cell types including xylem and fibre, a thick secondary wall comprised of cellulose, hemicellulose and lignin is deposited. Secondary cell walls provide a physical barrier that protects plants from pathogens, promotes tolerance to abiotic stresses and fortifies cells to withstand the forces associated with water transport and the physical weight of plant structures. Grasses have numerous cell wall features that are distinct from eudicots and other plants. Study of the model species *Brachypodium distachyon* as well as other grasses has revealed numerous features of the grass cell wall. These include the characterisation of xylosyl and arabinosyltransferases, a mixed-linkage glucan synthase and hydroxycinnamate acyltransferases. Perhaps the most fertile area for discovery has been the formation of lignins, including the identification of novel substrates and enzyme activities towards the synthesis of monolignols. Other enzymes function as polymerising agents or transferases that modify lignins and facilitate interactions with polysaccharides. The regulatory aspects of cell wall biosynthesis are largely overlapping with those of eudicots, but salient differences among species have been resolved that begin to identify the determinants that define grass cell walls.


\*These authors contributed equally to this work.

## I. Introduction to the secondary cell wall

The secondary plant cell wall provides mechanical strength that allows plants to stand upright, resist pest and pathogen invasion, and transport water over long distances. Both plants and humans have found this abundant matrix of cross-linked polymers useful as durable building material, with timber featuring in human construction around the world for generations. The secondary wall is distinct from other cell wall types in composition as well as the developmental timing and tissue types where it is deposited. Secondary walls form in thick layers, rich in cellulose, hemicelluloses and lignin. Cellulose microfibrils have a tensile strength rivaling steel, and form crystalline structures. Hemicelluloses include a variety of polysaccharides but, in grass secondary walls, these are mostly mixed-linkage glucans and heteroxylans, a defining aspect of this plant lineage. Finally, lignin is a recalcitrant and heterogeneous mixture of randomly polymerised phenolic monolignols that is interspersed and cross-linked with wall polysaccharide polymers. Lignification is a hallmark of secondary walls, and unique chemistry and synthesis of this polymer continues to be uncovered (Fig. 1).

Following cell expansion of cells surrounded by a primary wall, secondary walls are deposited in a highly specific spatio-temporal

manner in certain cell types over development. Unlike eudicots, grass stem growth is a result of iterative division and elongation events from stacked intercalary meristems called nodes (Esau, 1977; Langer, 1979). New cells generated from the node elongate, pushing up the nodes above with the final node transitioning to the flowering meristem. Thus, the internode regions are most mature at the bottom of the stem, while cells within an internode are most mature at the top of that region, just before the next node (Langer, 1979). Secondary wall deposition occurs between cell elongation and senescence, with cellulose, lignin and hemicellulose content increasing with maturity (Rancour *et al.*, 2012; Matos *et al.*, 2013; Kapp *et al.*, 2015). Grass stems account for the majority of secondary wall-forming sclerenchyma tissues. The interfascicular fibres develop thick secondary walls and provide mechanical strength for the upright stem. Grasses form discrete vasculature with the xylem and phloem contained by bundle sheath cells, unlike eudicots in which a cambium separates colateral xylem and phloem (Fig. 2). Depending on the species, stem vascular bundles can be arranged in peripheral rings or dispersed throughout the stem (Esau, 1977). Xylem develops strong secondary walls that can tolerate the high pressures caused by evapotranspiration. Phloem cells do not have secondary walls. Besides phloem, grass stem




**Fig. 1** General schematic of grass secondary cell wall matrix. The grass secondary cell wall matrix is made up of cellulose microfibrils, heteroxylans, mixed-linkage glucans, and lignins. (a) Generalised cartoon of grass secondary wall polymer interactions. (b) Schematic fine structure of the circled region in (a). Cellulose microfibrils consist of multiple, organised,  $\beta(1,4)$ -linked glucose chains. Mixed-linkage glucans are also glucose chains, but include  $\beta(1,3)$  linkages. Heteroxylan has a xylose backbone that is decorated with sugar and phenolic side chains of xylose, arabinose, glucuronic acid, and hydroxycinnamates (FA and *p*CA). These polysaccharides can be interwoven with lignins, branched phenolic polymers made of three main lignin units, syringyl (S), guaiacyl (G), and *p*-hydroxyphenyl (H). Lignins can also contain ferulic and *p*-coumaric acids.

parenchyma tissue can be found in the pith and in cortex pockets, which have been shown to function as carbon storage tissues during development (Jensen & Wilkerson, 2017). While this review focuses on secondary cell walls in *Brachypodium distachyon* and other grasses, wall synthesis has also been investigated using *B. distachyon* as a model system for callus tissue, young vegetative growth, and endosperm development (Christensen *et al.*, 2010; Guillon *et al.*, 2011; Liu *et al.*, 2016; Betekhtin *et al.*, 2018; Francin-Allami *et al.*, 2019).

## II. *Brachypodium distachyon*, a model grass system

*Brachypodium distachyon* is a model for cereal crops and temperate grasses because of its small stature, simple growth requirements, short life cycle, relatively small and sequenced genome, and close phylogenetic relation to those species (Scholthof *et al.*, 2018). *Brachypodium distachyon* has a ‘finished’ reference genome with the only ambiguity being the placement of some centromeric repeats ([https://phytozome-next.jgi.doe.gov/info/Bdistachyon\\_v3\\_1](https://phytozome-next.jgi.doe.gov/info/Bdistachyon_v3_1)). In addition, there is a growing atlas of gene expression profiles (Trabucco *et al.*, 2013; Sibout *et al.*, 2017; MacKinnon *et al.*, 2020). It is also remarkable in terms of the resources available for experimental molecular genetics. Genetic transformation

protocols are well developed; current efficiency makes *B. distachyon* a grass highly amenable to transformation (Bragg *et al.*, 2012). Mutant resources consist of 23 000 T-DNA mutants and 1200 sequenced chemical mutants (Bragg *et al.*, 2012; Granier *et al.*, 2015). Given that these mutations are more-or-less randomly distributed across the genome and chemical mutagenesis typically induces multiple mutations per mutant line, this large collection is likely to include loss-of-function mutations in the majority of *B. distachyon* genes and multiple nonsynonymous mutations in virtually every gene (Dalmis *et al.*, 2013). This latter category of mutations may be particularly interesting because it can help to elucidate the function of cell wall genes, as well as the importance of specific amino acids and protein domains, information that cannot be inferred from knockout mutants. A large natural variation population exists for *B. distachyon*, with sequenced genomes for many accessions. These resources have been applied in several studies on growth- and biomass-related traits (Lee *et al.*, 2012; Tyler *et al.*, 2014; Kapp *et al.*, 2015; Gordon *et al.*, 2017) Thus, *B. distachyon* is well positioned for gaining fundamental insights into cell wall biosynthesis (Coomey & Hazen, 2015). This knowledge can then be leveraged for agronomic gains in more experimentally recalcitrant grass species.



**Fig. 2** Transverse section of *Brachypodium distachyon* and *Arabidopsis thaliana* stems. Transverse stem cross-sections of *A. thaliana* (a) and *B. distachyon* (b) stained with phloroglucinol-HCl, a general stain for lignified tissues. Most eudicots, such as *A. thaliana*, have vascular bundles of xylem separated from phloem by cambium layers, and flanked by interfascicular fibres. In *B. distachyon*, the stem vascular bundles also contain xylem and phloem, but there is no cambial layer, and the vasculature is encased by a lignified bundle sheath layer of mestome cells and surrounded by interfascicular fibre cells. In both species, a cortex region of less lignified cells separates the interfascicular region from the epidermis. VB, vascular bundles; Ep, epidermis; Co, cortex; Ca, cambium area; IF, interfascicular fibres; Ph, phloem; Mes, mestome; Mx, metaxylem; Px, protoxylem; P, parenchyma. Bars, 100  $\mu\text{m}$ .

### III. Cellulose

Cellulose is perhaps the most abundant polymer in the world, found in the walls of every plant cell. It is made of (1,4)- $\beta$ -linked glucose monomers, and these glucan chains are synthesised at the plasma membrane by the cellulose synthase complex (Fig. 1). Extruded cellulose chains form organised microfibrils with crystalline structure; the degree of this organisation impacts wall mechanics, with greater crystallinity resulting in stiffer walls.

The cellulose synthase complex consists of multiple Cellulose Synthase A (CesA) subunits and associated proteins (Pear *et al.*, 1996; Polko & Kieber, 2019). *CesA* genes are a subclade of the cellulose synthase superfamily, along with the *Cellulose Synthase-like (Csl)* clades. Across plant species, seven major lineages have been identified in the *CesA* genes, which separate into the *CesAs* associated with primary or secondary wall synthesis (Little *et al.*, 2018). This distinction between primary and secondary wall synthesis is conserved across most vascular plants. In *B. distachyon*, *BdCesA4*, 7, and 8 have been shown to function in secondary wall synthesis, and these proteins are highly similar to those characterised in other species for secondary wall function, such as *Arabidopsis thaliana* and rice (*Oryza sativa*) (Handakumbura *et al.*, 2013). In *B. distachyon*, loss-of-function in the secondary *CesAs* results in reduced crystalline cellulose content, compromised wall integrity, and reduced plant growth (Handakumbura *et al.*, 2013; Petrik *et al.*, 2016). Interestingly, the secondary *CesA* lineage contains a Poacea-specific clade, which in *B. distachyon* is represented by *BdCesA10*. This *CesA10* group does not contain the canonical UDP-glucose binding motif (D,D,D,QxxRW) found in glucosyltransferases (Handakumbura *et al.*, 2013). While phylogenetic analysis clearly places these proteins in the *CesA* clade, it is not clear what role they play, if any, in cell wall synthesis.

Mutants in maize (*Zea mays*), barley (*Hordeum vulgare*), and rice with defects in cellulose synthesis have been identified through brittle stem phenotypes, aptly named brittle stalk, fragile stem, and brittle culm respectively (Tanaka *et al.*, 2003; Sindhu *et al.*, 2007; Burton *et al.*, 2010b; Kotake *et al.*, 2011). These mutants have been mapped both to genes encoding *CesAs* and other associated proteins, such as the COBRA-like family of glycosylphosphatidylinositol anchored proteins. While the precise function of these anchored proteins is not fully understood, they may play a role in properly orienting cellulose synthesis.

Cellulose synthase complex dynamics have been studied primarily in *A. thaliana*, but recent work in *B. distachyon* has added to our understanding of the conserved functions of this system. The complex moves along cortical microtubules, depositing cellulose microfibrils perpendicular to the axis of elongation (Paredes *et al.*, 2006). This has been observed in real time for primary *CesAs* in both *A. thaliana* and *B. distachyon*, which showed similar speeds in *B. distachyon* mesocotyl and root, as in *A. thaliana* hypocotyl. This motility was not affected by latrunculin B treatment, which destabilized actin filaments, but was dampened in both species when microtubules were disrupted (Liu *et al.*, 2017). Missense mutation in *Bdcesa1*, a primary wall cellulose synthase, showed reduced cellulose content and crystallinity, as do

*A. thaliana Atcesa1* mutants (Arioli *et al.*, 1998; Persson *et al.*, 2007; Brabham *et al.*, 2019). Unlike *Atcesa1* mutants, *Bdcesa1* did not show reduced plant height. Rather, the *Bdcesa1* mutant had more internodes, giving rise to a plant with normal height despite reduced cellular elongation from compromised cellulose synthesis (Brabham *et al.*, 2019). Overall, the process of cellulose biosynthesis appears to be somewhat conserved between eudicots and grasses.


### IV. Mixed-linkage glucans

One of the salient differences that defines grass secondary cell walls is the composition and utilization of noncellulosic polysaccharides. These can generally be thought of as pectins and hemicellulose, but discussion of these polymers is often better suited to classification by backbone structure (Scheller & Ulvskov, 2010; Atmodjo *et al.*, 2013). In eudicots, the predominant polysaccharide polymer after cellulose is xyloglucans, (1,4)- $\beta$ -linked glucose chains that contain numerous 1–6 xylose substitutions. The xylose side chains can be further decorated with other sugars such as galactose or fucose (Bauer *et al.*, 1973; Fry, 1989; Scheller & Ulvskov, 2010). In grasses, the role of xyloglucans is largely replaced by mixed-linkage glucans (MLGs) and heteroxylans.

Mixed-linkage glucans are, as their name suggests, (1,4)- $\beta$ -linked glucose chains that are interrupted with (1,3)- $\beta$  linkages (Figs 1, 3). (1,3)- $\beta$ -glucans are typically separated either by two or three (1,4)- $\beta$ -glucans, forming oligosaccharide units of  $\beta$ -cellotriosyl or  $\beta$ -cellotetraosyl (Fig. 3), although longer chains of (1,4)- $\beta$ -glucans are also observed (Bulone *et al.*, 2019). Almost no evidence of adjacent (1,3)- $\beta$ -glucan bonds has been found (Buliga *et al.*, 1986). These altered linkages result in the polymer having kinks and bends, unlike the linear glucan chains that form cellulose. As a result, MLG does not form crystalline structures. The relative amounts of  $\beta$ -cellotriosyl and  $\beta$ -cellotetraosyl units strongly relate to the solubility of the overall polymer and are expressed as ratios of degrees of polymerization of trisaccharides and tetrasaccharides (DP3 : DP4). Solubility of the polymer decreases at either end of the ratio spectrum. Longer stretches of either  $\beta$ -cellotriosyl or  $\beta$ -cellotetraosyl units increases the overall order of the polymer with more undisturbed regions of (1,4)- $\beta$ -glucan linkages, and thus decreases solubility. Greater solubility occurs with DP3 : DP4 ratios that range from 1 : 1 to 2.5 : 1 (Lazaridou & Biliaderis, 2007; Burton *et al.*, 2010a).

Mixed-linkage glucans were once thought to be unique to grass cell walls, but several examples have now been observed outside of the commelinid monocots, and indeed outside of green plants. Polysaccharides containing (1,3;1,4)- $\beta$ -glucans have been observed in green, red and brown algae, lichens, fungi, bryophytes, and the monophyletic genus *Equisetum* (Bulone *et al.*, 2019). Genomic data further support the idea that MLGs are not specific to the Poaceae, with enzymes capable of synthesizing (1,3;1,4)- $\beta$ -glucan linkages identified across monocots and in isolated cases in other species. MLG has been shown to be synthesized by members of the *CslF*, *CslH* and *CslJ* families (Bulone *et al.*, 2019). All three of these groups have co-evolved independently in monocots from sister *Csl* clades (Little *et al.*, 2018). Members of *CslF/HIJ* clades have been





**Fig. 3** Mixed-linkage glucan structure. (a) Fine structure of mixed-linkage glucan. Glucose monomers (yellow) linked by  $\beta(1,4)$  bonds (purple) are occasionally interrupted by  $\beta(1,3)$  linkages (pink). The  $\beta(1,3)$  bonds do not occur sequentially, but rather separate  $(1,4)$ - $\beta$ -glucans into  $\beta$ -cellobiosyl or  $\beta$ -cellobiosyl segments. The relative degree of  $\beta$ -cellobiosyl to  $\beta$ -cellobiosyl units relates to the solubility of the overall polymer.  $(1,3;1,4)$ - $\beta$ -Glucans are synthesised by Cellulose synthase-like F6, a Golgi membrane-bound protein with cytoplasmically active catalytic sites. (b) Miniature of Fig. 1(a) cell wall schematic highlighting the mixed-linkage glucan component.

shown to be capable of synthesizing  $(1,3;1,4)$ - $\beta$ -glucan when heterologously expressed, but it is not clear whether all of these groups are responsible for native MLG synthesis. By far the best characterized enzyme in MLG synthesis is CslF6, which has been studied in barley, wheat (*Triticum aestivum*), rice, maize and *B. distachyon* (Nemeth *et al.*, 2010; Vega-Sanchez *et al.*, 2012; Kim *et al.*, 2015, 2018). BdCslF6 protein is localized to the Golgi membrane, with an external catalytic domain (Kim *et al.*, 2015, 2018). Antibody detection of  $(1,3;1,4)$ - $\beta$ -glucan in maize also supports a Golgi-localized synthesis of MLG (Carpita & McCann, 2010). However, evidence in other grasses suggests that MLG synthesis occurs at the plasma membrane. In barley and wheat, antibody detection of MLG showed localization at the plasma membrane and cell wall, as did antibody detection of HvCslF6 and TaCslF6 (Trethewey & Harris, 2002; Trethewey *et al.*, 2005; Wilson *et al.*, 2006, 2015). The N-terminal region of the CslF6 protein in barley, maize, and sorghum (*Sorghum bicolor*) influences total MLG synthesis activity and the C-terminal region appears to influence the ratio of DP3:DP4 linkages (Jobling, 2015; Dimitroff *et al.*, 2016).

The evolution of MLG appears to have been followed by the evolution of hydrolytic enzymes specific to  $(1,3;1,4)$ - $\beta$ -glucan polymers (Høj & Fincher, 1995; Fincher, 2009). Both  $(1,4)$ - $\beta$ -glucan and  $(1,3)$ - $\beta$ -glucan endohydrolases exist across land plant lineages, capable of cleaving  $(1,4)$ - $\beta$ -glucan bonds in both cellulose and MLG. Specific  $(1,3;1,4)$ - $\beta$ -glucan endohydrolases have been well characterized in the metabolism of MLG, and analysis of their amino acid sequence and crystal structure shows strong similarity with barley  $(1,3)$ - $\beta$ -glucan endohydrolases, indicating that the ability to cleave  $(1,3;1,4)$ - $\beta$ -glucan polymers was achieved through

a modification of  $(1,3)$ - $\beta$ -glucan endohydrolase function (Varghese *et al.*, 1994).

The utility of increased MLG as a bioenergy source and the effect of increased MLG on wall content and plant health has been explored in studies on synthesis in barley and *A. thaliana* in which MLG was overexpressed. Excess MLG synthesis under constitutive promoters was detrimental to plant health, but tissue or developmentally specific promoters driving MLG synthesis resulted in plants with higher MLG content in grain or stem without such deleterious effects (Burton *et al.*, 2011; Vega-Sánchez *et al.*, 2015). In barley, MLG and starch levels have been shown to be inversely related in the developing coleoptile (Roulin *et al.*, 2002), and MLG levels have been shown to rise dynamically and fall over the course of development in vegetative tissue (Gibeau *et al.*, 2005). The grain cell walls of *B. distachyon* differ from those of cultivated cereals with exceptionally high levels of MLG and relatively lower starch levels (Guillon *et al.*, 2011; Opanowicz *et al.*, 2011; Trafford *et al.*, 2013; Burton & Fincher, 2014). This shift in carbon storage suggests that *B. distachyon* may rely on MLG to a greater extent than starch for endosperm carbon storage (Trafford *et al.*, 2013; Burton & Fincher, 2014). It has been suggested that MLG metabolism is enzymatically simpler than starch metabolism, requiring fewer enzymes in more available cellular spaces than the multistep, amyloplast specific process of starch metabolism (Roulin *et al.*, 2002; Burton & Fincher, 2012; Trafford *et al.*, 2013; Bulone *et al.*, 2019). While this has yet to be explored experimentally, it has been noted that a fast, alternative glucose storage pathway from  $(1,3;1,4)$ - $\beta$ -glucan metabolism may confer an advantage to the grasses, as evidenced by the development of this mechanism in a group with such widespread success.


## V. Grass heteroxylans

After glucans, xylans are the most abundant polysaccharide in plants. Although present across angiosperms, heteroxylans play a more prominent role in the grasses as the major hemicellulose (Scheller & Ulvskov, 2010). This class of polysaccharide is based on a (1,4)- $\beta$ -D-xylopyranosyl backbone, with side chains of arabinose, xylose, glucuronic acid, and hydroxycinnamates (Figs 1, 4). The nature and patterning of these side chains have major impacts on cell wall integrity, mediating xylan–cellulose and xylan–lignin polymer interactions (Simmons *et al.*, 2016; Martínez-Abad *et al.*, 2017). The  $\beta$ -(1,4)-xylan backbone has been shown to be synthesised in both eudicots and monocots by members of glycosyltransferase 43 (GT43) and GT47 family proteins. The *A. thaliana* irregular xylem mutants (*irx*) were some of the first identified xylan synthesis mutants, including *irx9*, *irx14* and *irx10*, all encoding GT43 and GT47 enzymes in wild-type plants (Brown *et al.*, 2005, 2009; Lee *et al.*, 2007; Peña *et al.*, 2007). In *B. distachyon*, recent work has shown that a member of the GT43 family is, in part, responsible for heteroxylan backbone synthesis. Genetic linkage mapping of saccharification rate in a recombinant inbred population identified a quantitative trait locus interval containing a *BdGT43A* orthologue of *A. thaliana* *IRX14* (Whitehead *et al.*, 2018). Allelic variation in *BdGT43A* between parental accessions Bd3-1 and Bd21 showed that the Bd3-1 allele encodes an alanine to threonine (A80T) shift that was associated with reduced Bd3-1 saccharification. Knockdown of *BdGT43A* resulted in reduced xylose, arabinose and ferulic acid deposition in stem tissue. Rice GT43 proteins have similarly been shown to mediate xylan synthesis, with *OsGT43A* and *OsGT43E* complementing *A. thaliana* *irx9* mutant phenotypes, and *OsGT43J* complementing *irx14* (Lee *et al.*, 2014).

The addition of side chains to the xylan backbone differentiates the various types of heteroxylans. In eudicots, glucuronoxylan is the most prevalent form, in which the side chain is formed by the addition of  $\alpha$ -(1,2)-GlcA side chains, sometimes amended with 4-

O-Me groups (Scheller & Ulvskov, 2010). Grass cell walls differ from those of eudicots in their abundance of arabinoxylans and glucuronoarabinoxylan. Arabinoxylans have monomer side chains of  $\alpha$ -(1,3)-Araf and  $\beta$ -(1,2)-Xylp, or dimer side chains of  $\alpha$ -(1,3)-Araf- $\alpha$ -(1,2)-Araf,  $\alpha$ -(1,3)-Araf- $\beta$ -(1,2)-Xylp, or  $\alpha$ -(1,3)-Araf ferulic acid. Glucuronoarabinoxylans contain the same side chains as arabinoxylans, but also include  $\alpha$ -(1,2)-GlcA-4-O-Me additions. Arabinoxylans are the more prevalent form found in endosperm cell walls, while glucuronoarabinoxylan is more common in vegetative tissue. The addition of these sugar side chains to heteroxylans is mediated by xylan arabinosyltransferases (XAT), which are members of the GT61 family. They function in the Golgi to add  $\alpha$ -(1,3)-Araf substitutions to the xylan backbone. Two XATs in wheat (*TaXAT1*, *TaXAT2*) and rice (*OsXAT2*, *OsXAT3*) have been characterised both natively and in heterologous systems for arabinosyltransferase activity (Anders *et al.*, 2012; Zhong *et al.*, 2018b). Other GT61 members possess xylosyltransferase activity. Rice xylosyl arabinosyl substitution of xylan 1 (*OsXAX1*) mediates the addition of xylose to arabinose units (*Xylp*-1,2- $\beta$ -Araf) (Chiniquy *et al.*, 2012), while rice xylan xylosyltransferase 1 (*OsXYXT1*) adds xylose sidechains to the xylan backbone (*Xylp*-1,2- $\beta$ -Xylp) (Zhong *et al.*, 2018b). While much of our understanding of heteroxylan synthesis comes from rice, some *B. distachyon* saccharification mutants identified from a sodium azide mutant population are candidates for characterising heteroxylan synthesis (Dalmais *et al.*, 2013). The *sac1* GT61 mutant in *B. distachyon* has a phenotype similar to the rice mutant *OsXAX1* (Marriott *et al.*, 2014). In *sac1*, plants have reduced xylose content, suggesting that the GT61 candidate, like *OsXAX*, mediates the incorporation of this saccharide component into the wall.

The presence of glucuronic acid (GlcA) side chains differentiates heteroxylans into glucuronoarabinoxylans and arabinoxylans. In *A. thaliana*, Glucuronic acid substitution of Xylan (AtGUX)-1 adds GlcA at evenly spaced intervals of 8–10 xylose residues, although greater spacing has been observed. AtGUX2 appears to preferentially



**Fig. 4** Grass heteroxylan structure. (a) Fine structure of heteroxylan and biosynthetic enzymes. The major noncellulosic polysaccharides in grasses are xylans. A xylose (blue) backbone is decorated with side chains of xylose, arabinose (green), glucuronic acid (teal), and ferulic acid (purple). The enzymes responsible for forming certain linkages on the heteroxylan polymer are depicted in either yellow, pink or both, having been characterised respectively in grass systems, eudicots or showing conserved function. (b) Miniature of Fig. 1(a) cell wall schematic highlighting the heteroxylan component.

add GlcA more frequently, at 5–7 residue intervals without regard for even spacing (Bromley *et al.*, 2013). The evenly spaced xylan regions form the major xylan domain, and the less organised GlcA spacing populates the minor domain. The major domain has been shown to interact with cellulose microfibrils, an interaction that is also mediated by xylan acetylation. Similar GUX function has yet to be observed in grasses, but presumably a mechanism for adding GlcA to heteroxytan exists. Additionally, 4-*O*-methylation of GlcA by AtGXMT (glucuronoxylan methyltransferase), a DUF579 protein, has been characterised in *A. thaliana*, but not in any grasses to date (Urbanowicz *et al.*, 2012). The addition of GlcA and its methylation have been implicated in eudicots in mediating xylan interaction with other wall polymers, and this phenomenon is ripe for investigation in grasses.

## VI. Xylan acetylation

Xylan acetylation has long been observed, but only recently has the role of these modifications been revealed. In *A. thaliana*, recent work has shown that acetylation pattern influences xyloans–cellulose interaction. Regularly spaced acetylation on every other xylose monomer in regions of the xylan backbone results in the polymer forming a two-fold helix that closely bonds with the hydrophilic side of cellulose microfibrils (Busse-Wicher *et al.*, 2014). The modification of xylan with acetate has strong implications for the solubility of the polymer, as well as the strength of xylan–cellulose interactions. Xylan-*O*-acetyltransferases (XOATs) are DUF231 family proteins, and carry out 2-*O*- and 3-*O*-monoacetylation and 2,3-di-*O*-acetylation (Fig. 4). In *A. thaliana*, nine XOATs have been identified and genetically characterised, including the Trichome Birefringence protein, TBR-like proteins and ESKIMO1 (Zhong *et al.*, 2017). In grasses, there has been an expansion of the DUF231 XOATs, with rice containing 14 members. *OsXOAT1* and *OsXOAT7* complement the *A. thaliana esk1* xylan acetylation mutant, and all 14 rice XOATs can acetylate xylohexose *in vitro* (Zhong *et al.*, 2018a).

While the degree of xylan acetylation has been shown to play a critical role in wall integrity, evidence of deacetylation activity has not yet been shown in eudicots. However, *rice brittle sheath 1* (*OsBS1*) encodes a GDSL lipase/esterase that functions as an acetyltransferase in the Golgi, removing acetyl groups from xyloans (Zhang *et al.*, 2017). Mutation in *OsBS1* results in greater 2-*O*- and 3-*O*-acetylation, which compromises secondary wall patterning and integrity.

## VII. Lignins


Lignins are large phenolic polymers mainly deposited in the primary and the secondary cell wall of xylem and sclerenchyma cells. These polymers provide the hydrophobicity and mechanical properties necessary for the development of land plant vasculature. Lignins embed polysaccharides in the cell wall and are a major barrier for biomass usage such as saccharification for biofuel production (Marriott *et al.*, 2014). Unlike other wall polymers, lignins contain many types of interunit bonds (aryl beta-aryl ether, phenyl coumaran, resinol, biphenyl) randomly formed during

polymerisation, some being more (C–C) or less (C–O–C) resistant to degradation (Mnich *et al.*, 2020). Consequently, lignin structure is not predictable, although the abundance of each monomer seems to influence the occurrence of certain linkages (Stewart *et al.*, 2009).

Lignins are synthesised from three monolignols, *p*-coumaryl alcohol, coniferyl alcohol and sinapyl alcohol, that differ by their degree of methoxylation. Once incorporated into lignin polymers, these phenolics give rise to *p*-hydroxyphenyl (H), guaiacyl (G), and syringyl (S) units, respectively. In *B. distachyon* stems, lignin content accounts for 18–25% of the dry cell wall residue and, in the wild-type Bd21-3 accession, stem lignin is comprised of about 62% S, 34% G, and 4% H units (Bouvier d'Yvoire *et al.*, 2013; Trabucco *et al.*, 2013). Within the grasses, *B. distachyon* has one of the highest proportions of S units reported (Clarke *et al.*, 1933; Méchin *et al.*, 2014; Herbaut *et al.*, 2018).

Lignin biosynthesis results from a branch of the phenylpropanoid pathway and has long been thought to rely on the aromatic amino acid L-phenylalanine (L-Phe) as a starting substrate (Fig. 5). The standard convention in most studied plant systems has been that L-Phe is first deaminated by phenylalanine ammonia lyase (PAL), yielding cinnamate, which is then C4-hydroxylated by coumarate-4-hydroxylase (C4H) to make coumarate. Coumarate is a common branch point for all three main monolignols. However, this conventional pathway has recently been challenged by work in *B. distachyon* demonstrating that L-tyrosine (L-Tyr) can also serve as an initial substrate for lignin synthesis as it already contains a C4 hydroxylation. Indeed, tyrosine ammonia lyase (TAL) activity in grasses (Higuchi *et al.*, 1967) suggests that C4H activity can be bypassed to produce coumarate (Fig. 5). In grasses, PAL and TAL activities are controlled by the same protein, but clear evidence for a genuine phenylalanine tyrosine ammonia lyase (PTAL) activity in the phenolic pathway was poorly documented until recently. In plants that expressed a *BdPAL* RNAi hairpin construct to knock down expression of multiple *BdPAL* genes, both PAL and TAL activities were affected and plants contained 43% less lignin (Cass *et al.*, 2015; Barros *et al.*, 2016). Only one predicted PTAL (PTAL1) was identified in this family and nearly half of the total lignin deposited in *B. distachyon* occurs via TAL activity (Barros *et al.*, 2016). Interestingly, *BdPTAL1* is mainly involved in the biosynthesis of S units and cell wall linked coumarates, with less effect on G units as revealed by plants fed with C13-labelled L-Phe or L-Tyr. A biological role for PTAL has only been shown in *B. distachyon* to date, but putative orthologues to *BdPTAL1* have been identified in several other grasses (Barros *et al.*, 2016). Further characterisation is needed to confirm whether this alternate initiation of lignin synthesis is shared broadly amongst grasses, or indeed present in other groups.

Other recent discoveries are further changing our understanding of lignin biosynthesis in grasses. Very recently, (Barros *et al.*, 2019) proposed that a cytosolic ascorbate peroxidase with genuine 4-coumarate 3-hydroxylase (C3H) activity oxidises coumarate into caffeate in the phenylpropanoid pathway. Decreased expression of this novel C3H in *B. distachyon* results in significantly reduced lignin content and structure. This 'acid' route to caffeic acid and thus to caffeoyl CoA through the activity of 4-hydroxycinnamate



**Fig. 5** Alternative routes to lignin biosynthesis. (a) In some grasses, phenylalanine tyrosine ammonia lyase (PTAL) can bypass cinnamic acid 4-hydroxylase (C4H) activity and thus directly produce *p*-coumarate from tyrosine. An ascorbate peroxidase-like gene with coumarate 3-hydroxylase (C3H) activity can directly synthesise caffeate from coumarate. Red arrows illustrate a grass-specific lignin pathway unique from other lignin pathways identified in most studied eudicots. Green arrows show routes potentially bypassed or absent in *Brachypodium distachyon*, such as caffeoyl shikimate esterase (CSE). Blue arrows are paths believed to be common to eudicots and grasses. Asterisks highlight enzymes characterised in *B. distachyon*. The mechanism for monolignol export across the plasma membrane remains unclear. Phenylalanine ammonia lyase (PAL), 4-hydroxycinnamoyl CoA ligase (4CL), hydroxycinnamoyl CoA:shikimate hydroxycinnamoyl transferase (HCT), *p*-coumaroyl shikimate 3'-hydroxylase (C3'H), caffeoyl CoA O-methyltransferase (CCoAOMT), hydroxycinnamoyl CoA reductase (CCR), ferulic acid 5-hydroxylase (F5H), caffeic acid/5-hydroxyferulic acid O-methyltransferase (COMT), cinnamyl alcohol dehydrogenase (CAD), laccase (LAC), peroxidase (PRX). R- in the monolignol hydroxycinnamics could be substituted by a hydrogen or methyl group. (b) Miniature of Fig. 1(a) cell wall schematic highlighting the lignin component.



CoA ligase (4CL) would be complementary to the C3'H pathway in which 4CL, 4-hydroxycinnamoyl CoA:shikimate/quinic acid hydroxycinnamoyltransferase (HCT), 4-coumaroyl shikimate/quinic acid 3'-hydroxylase (C3'H), function sequentially to convert coumarate to caffeoyl CoA (Fig. 5). Feruloyl-CoA produced by the methoxylation of caffeoyl CoA by caffeoyl CoA *O*-methyltransferase (CCoAOMT) is a substrate of cinnamoyl-CoA reductase (CCR). The proposition of an alternate 'acid' route to monolignol synthesis is not new, but the discovery of a cytosolic ascorbate peroxidase with 4-coumarate 3-hydroxylase activity *in planta* was lacking until now. Interestingly, Barros *et al.* (2019) showed this is not unique to grasses, as the null allele of the *A. thaliana* C3H orthologue is lethal. The lignin pathway that involves the membrane-bound C3'H also plays a critical role in grass lignin synthesis. Indeed, C3'H-knockout rice mutants were severely affected in their development and displayed typical C3'H phenotypes with lignins largely enriched in H units at the expense of G and S units (Takeda *et al.*, 2018). Interestingly, caffeoyl shikimate esterase (CSE) activity was not detected in *B. distachyon* stem crude extract and this result is supported by the absence of close orthologues of *AtCSE* in *B. distachyon* (Ha *et al.*, 2016). CCR is a cornerstone step to monolignol biosynthesis. CCR activity converts CoA-conjugated intermediaries into the aldehyde precursors of monolignols. While *ccr* mutants with decreased lignin content and increased monolignol conjugates were studied in maize there are no reports on cell wall properties of CCR-deficient lines in *B. distachyon* (Tamasloukht *et al.*, 2011; Cass *et al.*, 2015).

The last enzyme in the monolignol pathway, cinnamyl alcohol dehydrogenase (CAD), reduces cinnamaldehyde into alcohols. Mutants and transgenics lines affected in CAD have been well characterised in *B. distachyon* (Bouvier d'Yvoire *et al.*, 2013; Trabucco *et al.*, 2013). Lignin content of *Bdcad1* mutants was drastically enriched in aryl  $\beta$ -aryl ether and diaryl ether-coupled S units, as well as resistant interunit bonds and free phenolic groups, a result previously observed in maize and sorghum *brown-midrib* mutants (Pillonel *et al.*, 1991; Barriere *et al.*, 2004). By contrast, there was little increase in coniferaldehyde-end groups in the *Bdcad1*, suggesting that another CAD gene specific to coniferyl alcohol is involved in lignification. As observed in CAD-deficient eudicot plants, sinapic acid esters linked to the cell wall were detected in *Bdcad1* (Bouvier d'Yvoire *et al.*, 2013).

As stated at an earlier point, *B. distachyon* lignin is relatively enriched in S units. Their precursor, sinapyl alcohol, is produced through the C5 hydroxylation of coniferaldehyde by the P450 enzyme ferulate-5-hydroxylase (F5H) and methoxylation by caffeoyl-*O*-methyltransferase (COMT). When *F5H* was overexpressed in *B. distachyon*, cell wall analysis revealed an average increase of 25% in the content of S units in these lines, leading to an increase in the S/G ratio from 2.3 in wild-type to 8.1, with a modest increase of 5-hydroxy-guaiacyl units (5-OH-G UNIT) and 30% higher saccharification yield (Sibout *et al.*, 2017). Several *B. distachyon* mutants affected in COMT activity were identified in a sodium azide-induced mutant collection by TILLING (Dalmais *et al.*, 2013). As observed in maize *comt* mutants, *B. distachyon* mutants showed the accumulation of 5-OH-G units in their lignin and significantly altered lignin

content (Piquemal *et al.*, 2002; Bouvier d'Yvoire *et al.*, 2013; Dalmais *et al.*, 2013; Trabucco *et al.*, 2013).

Once exported into the apoplast by an unknown mechanism, monolignols are oxidised by peroxidases and/or laccases (Vanholme *et al.*, 2012; Wang *et al.*, 2013; Perkins *et al.*, 2019; Vermaas *et al.*, 2019). Laccases are multicopper oxidases that use oxygen as an electron acceptor, while peroxidases use H<sub>2</sub>O<sub>2</sub>. Once oxidised, the monolignols radically polymerise into the branched lignin polymer with multiple bond types resulting from the various positions of the oxygen radical on the monolignol subunit. There are 17 laccases in *A. thaliana* and 29 in *B. distachyon* (Berthet *et al.*, 2011; Le Bris *et al.*, 2019). *Brachypodium distachyon* LACCASE 5 and 8 were identified as orthologues of *AtLAC17* and were shown to be responsible for lignification in interfascicular fibres (Wang *et al.*, 2015; Le Bris *et al.*, 2019). A laccase gene from sugarcane (*SofLAC*) also genetically complemented an *A. thaliana lac17* mutant (Cesarino *et al.*, 2013). Lignin content decreased by 30% in the double *lac5lac8* mutant, and saccharification increased by 140% compared with the wild-type. Lignin deposition was less affected in vascular bundles compared with fibres, suggesting that different laccases or peroxidases are recruited for lignification of these tissues.

## VIII. Hydroxycinnamic acids

The presence of hydroxycinnamic acids, namely ferulic acid (FA) and *p*-coumaric acid (*pCA*), in the cell wall is a defining feature of grass secondary cell walls (Ralph *et al.*, 1994; Hatfield *et al.*, 2009). FA is predominantly linked to heteroxylan through an ester bond. The oxidation of FA in the cell wall, probably by peroxidases, generates esterified dehydrodiferulates which serve as linkages between two arabinoxylan polymers. In lignified tissues, xylan-esterified ferulates can be etherified to G units of lignin and thus serve as a covalent linkage between hemicelluloses and lignins (Hatfield *et al.*, 2016; Lapierre *et al.*, 2019). An esterified ferulate on arabinoxylan is considered as a nucleation site of lignification in grasses and thus an important mechanism for cell wall reinforcement (Ralph *et al.*, 1995, 1998). *pCA* is esterified on arabinoxylan to a lesser extent than FA and tends to be found esterified to S units in *B. distachyon* lignins. Plant-specific acyl-CoA-dependent acyltransferases of the BAHD (BEAT, AHCT, HCBT, and DAT) family are the enzymes responsible for the acylation of the arabinose side chains of heteroxylans and monolignols with hydroxycinnamates (D'Auria, 2006; Mitchell *et al.*, 2007). An expanded grass-specific BAHD clade (also called the 'Mitchell clade') was identified by bioinformatic analysis in rice as candidates for hydroxycinnamate transfer (D'Auria, 2006; Mitchell *et al.*, 2007; Bartley *et al.*, 2013). Consequently, BAHD enzymes with feruloyl transferase activity were first explored in rice and have also been investigated in *B. distachyon* (Piston *et al.*, 2010; Bartley *et al.*, 2013). BAHD01 in *B. distachyon* and *Setaria viridis* appear to be involved in feruloylation of arabinoxylans (de Souza *et al.*, 2018). Downregulation of *SvBAHD01* significantly reduced FA on arabinoxylan, with an increase in *pCA*-Arafacylation and no substantial change in lignin content while in *B. distachyon* only a moderate decrease in FA-arabinoxylan was observed (de Souza *et al.*, 2018). Interestingly,

*BdBAHD01* downregulation lines showed increased saccharification efficiency, despite unchanged lignin content, highlighting the role of FA in maintaining cell wall integrity. Overexpression of *BdBAHD05* (also called *BdAT1*) caused a moderate increase in FA content and downregulation showed a moderate decrease (Buana-fina *et al.*, 2016; de Souza *et al.*, 2018). Analysis in sugarcane revealed six *BAHD* genes, one of which is homologous to *SvBAHD01*, and downregulation of *SacBAHD01* similarly reduced stem FA content and increased biomass digestibility (de Souza *et al.*, 2019).

FA acylated monolignols were detected in several species, including willow (*Salix* sp.) and poplar (*Populus trichocarpa*), although in much lower amounts than *pCA* acylated monolignols in grasses (Karlen *et al.*, 2016). In rice, feruloyl monolignol transferase (*OsFMT*) was identified through homology with other *BAHD* acyltransferases that act on monolignols (Wilkerson *et al.*, 2014; Karlen *et al.*, 2016). *OsFMT* overexpression resulted in higher levels of FA on lignin. FA from heteroxylan, released through mild alkaline hydrolysis, was unchanged by altered *OsFMT* expression. Furthermore, there was no change in the levels of *pCA* acylated monolignols, suggesting specificity of this enzyme for monolignol feruloylation.

In *B. distachyon*, *p*-coumaroyl-CoA:monolignol transferase (*PMT*) acylates lignin with *pCA*, but not heteroxylan (Petrik *et al.*, 2014). While *OsPMT* has a high affinity for coumaroyl alcohol *in vitro*, *BdPMT* preferentially acylates sinapyl alcohol with *pCA in planta* (Withers *et al.*, 2012; Sibout *et al.*, 2016). Lines overexpressing *BdPMT* showed lower total lignin despite an increase of *pCA* content (Petrik *et al.*, 2014). This may be a consequence of redirecting *p*-coumaric acid CoA for acylation rather than monolignol synthesis, or the inhibition of the monolignol polymerisation by excessive *p*-coumaroylation (Sibout *et al.*, 2016). Interestingly, when *BdPMT* was overexpressed in *A. thaliana*, which does not natively produce *pCA* acylated lignin, a significant amount of *pCA* was found on lignins (Sibout *et al.*, 2016). More surprising, when *BdPMT* was expressed under a specific *C4H* promoter in a *ccr*-deficient *A. thaliana* mutant background, lignin was esterified with both *pCA* and FA. Mutants in *CCR* accumulate high levels of feruloyl-CoA, and *BdPMT* activity in this mutant suggests that not only is *BdPMT* functional in eudicots, but it is also able to use feruloyl-CoA as a substrate when available in sufficient quantities (Withers *et al.*, 2012; Sibout *et al.*, 2016). In maize, *ZmPMT* loss-of-function lines had less *pCA* and modified lignin structure, but not reduced total lignin content (Marita *et al.*, 2014). Overall, *BAHD* proteins have a related set of functions in decorating cell wall components; feruloylation of arabinoxylan (*BAHD01*), feruloylation of lignins (*FMT*), and coumaroylation of lignins (*PMT*). An enzyme responsible for the coumaroylation of arabinoxylan remains to be discovered.

## IX. Tricin

As evidenced by their *pCA* and FA content, grasses are remarkable in their capacity to incorporate phenolic compounds other than the typical coumaryl, coniferyl and sinapyl alcohols into lignin. Tricin, an *O*-methylated flavone, was first characterised in wheat

straw lignin (del Río *et al.*, 2012). Tricin is incorporated into grass lignin in varying amounts across grass species, with oat (*Avena sativa*), wheat and *B. distachyon* straw being particularly enriched in this compound (Lan *et al.*, 2016). Tricin is incorporated in lignin polymers via 4'-*O*- $\beta$  coupling (Lan *et al.*, 2018). Biomimetic radical coupling reactions give evidence that tricin may serve as a possible nucleation site for lignification, as has been suggested for ferulate (Ralph *et al.*, 1995, 1998; Lan *et al.*, 2015). Tricin and monolignols come from two different branches of the phenylpropanoid pathway, and consequently their synthesis shares some common enzymes. This is particularly true for enzymes involved in the metabolic flux upstream of *p*-coumaric acid synthesis. *CHALCONE SYNTHASE*, a pivotal enzyme for flavonoids production, uses malonyl-CoA and *p*-coumaroyl-CoA as substrates. Silencing this enzyme in maize resulted in strongly reduced levels of apigenin-related and tricin-related flavonoids, and also strongly reduced incorporation of tricin into the lignin polymer (Eloy *et al.*, 2017). The effect of the flavonoid pathway on the production of cell wall tricin content was also demonstrated in rice (Lam *et al.*, 2017, 2019). It is also possible that some of the cell wall changes observed in *BdPMT* overexpression lines may stem from the depletion of *p*-coumaroyl-CoA pool, as both chalcone synthase and *PMT* act on this substrate. *O*-methyltransferases involved in the *O*-methylation of 5-hydroxy-coniferinaldehyde to produce sinapyl alcohol were also shown to be involved in the methylation of tricin in rice, maize, and sorghum (Eudes *et al.*, 2017; Fornalé *et al.*, 2017; Lam *et al.*, 2019). The bi-functionality of *COMT* in the lignin and flavonoid pathways is not unexpected, as a *COMT* involved in lignification of *A. thaliana* stems also *O*-methylates isorhamnetin, a flavonoid structurally similar to tricin (Do *et al.*, 2007). There is now abundant evidence that other molecules, called 'nontraditional monomers' like tricin or hydroxycinnamic acids, can be incorporated into the lignin polymer (del Río *et al.*, 2018; Vanholme *et al.*, 2019). The biological role of these novel lignin components remains to be determined.

## X. Silicon

Poaceae accumulate high quantities of silicon in the cell wall of their shoots. This phenomenon is particularly marked in rice (Ma & Yamaji, 2006). The main role of silicon is to provide plant resistance to many biotic and abiotic stresses (Hattori *et al.*, 2005; Deshmukh *et al.*, 2017). However, silicon may interact with polysaccharides, which consequently affect plant biomass processing in biorefineries (Perry & Lu, 1992; Kido *et al.*, 2015). For biofuel production, there is a trade-off between soil amendment with silicon that can increase polysaccharide yield with a negative effect on the conversion of biomass into biofuels (Głazowska *et al.*, 2018b). Silicon content in rice and maize can be modulated by changing the expression of silicon transporters (Ma *et al.*, 2007; Mitani-Ueno *et al.*, 2016; Bokor *et al.*, 2017). The analysis of different silicon transporter mutants showed that silicon availability may impact the morphology and patterning of stem and leaf macrohairs (Głazowska *et al.*, 2018b). The *Bd low silicon 1* (*Bdlsi1*) mutant is impaired in silicon transporter function and has reduced

silicon uptake, with 93% less silicon present in the shoot. Mixed-linkage glucan content is drastically modified in *Bdlsi1* (Kido *et al.*, 2015; Głazowska *et al.*, 2018a). This result is in agreement with previous studies suggesting that (1,3;1,4)- $\beta$ -glucan is involved in silicon-dependent strengthening of the rice cell wall (Kido *et al.*, 2015). The *Bdlsi1* mutant also displayed an altered degree and pattern of homogalacturonan methyl esterification. Despite the relatively low amount of pectins found in grasses, this change in homogalacturonan represents a significant alteration to the wall matrix. Lastly, *Bdlsi1* mutant FA extratability was lower with only minor changes in lignin content (Kido *et al.*, 2015; Głazowska *et al.*, 2018a). These data highlight the important role silicon plays in cell wall integrity in *B. distachyon* and grasses in general, and presents interesting avenues for further study.

## XI. Transcriptional regulation of secondary cell wall thickening


Canonical transcription factors that directly bind DNA play a prominent role in the regulation of plant secondary cell wall thickening. The *cis*-regulatory regions of genes associated with cellulose, hemicellulose and lignin biosynthesis interact directly with numerous MYB and NAC family transcription factors (Fig. 6; Nakano *et al.*, 2015). Many of the R2R3-MYB protein family subgroups appear to bind a similar sequence motif, the AC element, also known as the M46RE (MYB46 responsive *cis*-regulatory element) and the SMRE (secondary wall MYB-responsive element) (Kim *et al.*, 2012; Zhong & Ye, 2012; Zhao & Bartley, 2014; Handakumbura *et al.*, 2018). In *A. thaliana*, *AtMYB46* and the close paralogue *AtMYB86* activate the expression of cellulose, hemicellulose and lignin biosynthetic genes, as well as other MYBs capable of activating secondary cell wall-related genes (Zhong *et al.*, 2007; Zhong & Ye, 2007). Some of the downstream MYB activators, among them *AtMYB58/63* and *AtMYB42/85*, activate only lignin genes (Rao & Dixon, 2018; Zhang J *et al.*, 2018). However, in sorghum, rice, and switchgrass (*Panicum virgatum*), ectopic expression of *OsMYB58/63*, *PvMYB58/63*, and the sorghum orthologue *SbMYB60* results in the activation of cellulose and hemicellulose genes as well as lignin (Noda *et al.*, 2015; Scully *et al.*, 2016; Rao *et al.*, 2019). A potential orthologue to *OsMYB42/85*, *ZmMYB167*, was overexpressed in maize and heterologously in *B. distachyon* to similar effect (Bhatia *et al.*, 2019). Similar functions have been resolved for the *A. thaliana* and rice orthologues *AtMYB61* and *OsMYB61* as well as *AtMYB103* and *OsMYB103* (Hirano *et al.*, 2013; Huang *et al.*, 2015; O'Malley *et al.*, 2016; Zhao *et al.*, 2019). These downstream MYBs bind the AC element and activate both lignin and wall polysaccharide biosynthesis genes. Overall, there are few distinctions in the transcription factor targets for these genes between grasses and *A. thaliana*. Those that have been observed may be the outcome of low-resolution experimental designs that sample one tissue type at one time point for a limited number of outputs.

The expression of cell wall-associated genes is often highly correlated (Brown *et al.*, 2005; Persson *et al.*, 2005). Coexpression analysis of a *B. distachyon* gene expression atlas resolved a cluster of 96 genes that is enriched for cell wall biosynthetic processes with

numerous cellulose-, hemicellulose-, and lignin-associated genes (Sibout *et al.*, 2017). Among the identified genes, there are two primary and two secondary wall *CESAs*, as well as *COBRA*, *KORRIGAN*, *CSII*, *CSLF2*, numerous glycosyltransferases and glycosylhydrolases, fasciclin-like family and numerous lignin-associated genes. The MYB transcription factor *SECONDARY WALL ASSOCIATED MYB 1 (SWAMI)* is one of two canonical transcription factors that are part of the wall gene enriched cluster, making it a candidate for a regulator of genes in the cluster (Fig. 6). Similarly, analysis of *B. distachyon* leaf, root, and stem microarray gene expression data identified *SWAMI1/2/3* and MYBs that are part of six other prominent subgroups orthologous to *AtMYB46/83*, *AtMYB103*, *AtMYB58/63*, *AtMYB52/54*, *AtMYB42/85*, and *AtMYB4/32* that are highly correlated with secondary *CESA* and lignin biosynthetic gene transcriptional targets (Supporting Information Table S1; Handakumbura *et al.*, 2018). Interestingly, the *SWAMI* gene and its two closest homologues, *SWAM2* and *SWAM3*, are conspicuously absent from genomes in the *A. thaliana* family Brassicaceae but present in other eudicots and monocots (Handakumbura *et al.*, 2018). Like the other described secondary cell wall regulating R2R3-MYBs, *SWAMI* interacts with the AC element and is an activator of secondary cell wall genes. Based on amino acid similarity with characterised genes in other systems and their expression pattern, all of the *B. distachyon* identified MYBs are excellent candidates for a role in cell wall biosynthesis.

The same promoters that interact with the secondary cell wall regulating MYB transcription factors often interact with NAC transcription factors, collectively referred to as the *SECONDARY WALL NACs (SWN)* or the *VND*, *NST/SND*, *SMB* related (*VNS*) (Ohtani *et al.*, 2011; Zhong *et al.*, 2011). This group of proteins is generally classified into four clades, all binding the similarly named *VNS* element in *in vitro* assays (O'Malley *et al.*, 2016; Olins *et al.*, 2018), which is consistent with independently identified *TERE* and *SNBE* binding sites for the same proteins (Pyo *et al.*, 2007; Valdivia *et al.*, 2013). In *A. thaliana*, three of the clades that include the *VASCULAR-RELATED NAC-DOMAINS (VNDs)*, activate cell wall thickening directly and by activating the previously described downstream MYBs (Kubo *et al.*, 2005). The *VNDs* can induce vascular cells differentiation, induce further thickening, and initiate programmed cell death (Kubo *et al.*, 2005; Zhong *et al.*, 2008). They function in xylem rather than fibres where thickening is activated by the clade IV *NACs*: *NAC SECONDARY WALL THICKENING FACTOR 1 (NST1)*, *NST2*, and *NST3* (also known as *SECONDARY WALL-ASSOCIATED NAC-DOMAIN PROTEIN 1 (SND1)*) (Zhong *et al.*, 2006; Mitsuda *et al.*, 2007). Uniquely, programmed cell death is not activated by clade IV *NACs*. Such cell type specific functions have not been resolved in grasses. The function of the *SWNs* is well conserved between *A. thaliana* and grasses where grass genes can complement mutants in *A. thaliana* (Zhong *et al.*, 2011, 2015; Rao *et al.*, 2019). In *B. distachyon*, members of all four clades induced the formation of secondary walls when ectopically expressed in tobacco leaves and the *VND*-type *SWNs* also activated programmed cell death (Valdivia *et al.*, 2013). Together with the MYBs, the *NACs* form feed-forward loops (Nakano *et al.*, 2015; Taylor-Teeple *et al.*, 2015). In general, all of the transcription factor proteins can bind to





**Fig. 6** Transcriptional regulation of secondary cell wall deposition. Secondary cell wall transcription regulatory network in *Arabidopsis thaliana* (a) and grasses (b). Bean shape indicates DNA binding transcription factors. Ovals indicate protein interactors. Circles are hormones. Orthology between *A. thaliana* and grasses is denoted by colour. Blue, red and green arrows indicate regulation of cellulose, lignin and hemicellulose, respectively. Arrows indicate activation and bars repression. Grey shaded items have not been described in both systems.



genes that encode cell wall structural enzymes and they also have the function of activating other activating transcription factors.

The finger-like protuberance formed by a zinc-finger protein domain can interact with DNA, RNA and proteins. Tandem CCCH zinc-finger (TZF) proteins modulate gene expression transcriptionally by interactions with DNA or posttranscriptionally by interactions with mRNA (Bogamuwa & Jang, 2014). *Arabidopsis thaliana* *C3H14* is a direct activator of lignin, cellulose and hemicellulose biosynthesis genes and may be a repressor of MYB cell wall activators (Ko *et al.*, 2009; Kim *et al.*, 2012). Conversely, the *INDETERMINATE* family C2H2-type zinc-finger transcription factor in rice, OsIDD2, interacts with the ID motif to repress the expression of lignin-associated genes (Huang *et al.*, 2018). Analysis of mutants and transgenic plants suggests that OsIDD2 directly represses lignin-associated gene expression and indirectly secondary wall *CESA* genes. By yet another possible mechanism of gene regulation, the rice TZF protein ILA1-interacting protein 4 (IIP4) functions as a repressor of secondary wall thickening through protein–protein interaction with OsSWN2 (NAC29) and SWN3 (NAC31) (Zhang D *et al.*, 2018). The association with the SWNs is attenuated by phosphorylation of IIP4 protein, which results in translocation to the cytosol. Thus, zinc-finger proteins influence the thickening of grass secondary walls through multiple mechanisms.

While the regulatory network is dominated in number by MYB and NAC transcription activators, several repressors have also been described, namely Three Amino acid Loop Extension (TALE), zinc-fingers, HD-ZIP III, WRKY, LATERAL ORGAN BOUNDARY (LBDs) and some MYBs. Among the many types of cells that do not have secondary cell walls are pith, which reside inside the stem and among the cells in the plant with the thickest walls. A WRKY transcription factor, *WRKY12*, is a repressor of wall thickening in pith and other cells. It can directly bind the promoters of *AtNST2* and poplar *C4H* and broadly repress wall thickening in *A. thaliana*, poplar, and switchgrass (Wang *et al.*, 2010; Yang *et al.*, 2016; Rao *et al.*, 2019). The five class III HD-ZIPs in *A. thaliana* (*REVOLUTA*, *PHABULOSA*, *PHAVOLUTA*, *CORONA*, and *HB8*) and some orthologues in poplar have been shown to play a role in cambium cell initiation and vascular bundle organisation (Floyd & Bowman, 2006). However, to our knowledge, there are no reports describing a function for this group of genes in grasses. Several LBD family transcription factors, *AtLBD15/18/30*, can activate the expression of *AtVND7* and induce wall thickening and differentiation into tracheary cells (Soyano *et al.*, 2008; Ohashi-Ito *et al.*, 2018). *AtLBD29*, conversely, is a repressor of stem secondary wall thickening and is activated by the phytohormone auxin (Lee *et al.*, 2019). Repression is also supplied by the MYB G4 clade and are the best characterised in grasses. These include ZmMYB11/31/42, PvMYB4/32, and OsMYB108, which are orthologous to AtMYB4/32 (Zhao & Bartley, 2014; Rao & Dixon, 2018; Miyamoto *et al.*, 2019). These were first described in a grass as direct repressors of lignin gene expression (Fornalé *et al.*, 2006; Sonbol *et al.*, 2009). In switchgrass, PvMYB4 is a direct repressor of lignin-associated genes (Shen *et al.*, 2012; Rao *et al.*, 2019). The MYB31/42 MYBs in sorghum, rice and maize directly bind to the *cis*-regulatory regions of various lignin biosynthetic gene, but there

appears to be variation in phenylpropanoid gene expression and protein–DNA interactions across species (Agarwal *et al.*, 2016). Wounding-induced lignification occurs in maize by degradation of ZmMYB11/31/42 protein and a protein interacting partner ZML2 (Vélez-Bermúdez *et al.*, 2015). Thus, wounding and the subsequent activation of MeJA signalling will remove MYB G4 clade repression in maize and induce lignin gene expression. The repressing MYB G4 clade interacts with AC-like sequence motifs, similar to the wall activating MYBs (Fornalé *et al.*, 2010; Shen *et al.*, 2012; Agarwal *et al.*, 2016). The exact targets in the phenylpropanoid pathway vary across system and study, which suggests that there may be some transcription factor subfunctionalisation.

Members of two different classes of the TALE superfamily, KNOX and BEL, have been shown to regulate secondary wall synthesis. The class II KNOX gene *KNOTTED OF ARABIDOPSIS THALIANA 7* (*AtKNAT7*) was initially identified as an irregular xylem mutant (*ix11*) (Brown *et al.*, 2005). *KNAT7* orthologues are generally described in the literature as repressors and, while there is substantial evidence for this, there are also some outstanding issues raised by data indicating a role as an activator of wall deposition. *Atknat7* mutants have thicker interfascicular fibre walls, as expected for a repressor mutant, but this mutant also shows collapsed xylem (Brown *et al.*, 2005; Li *et al.*, 2012). *Atknat7* mutants have greater lignin content, but reduced xylan, suggesting that *AtKNAT7* may differentially regulate aspects of wall polymer synthesis. In conflicting reports, one group has shown xylan biosynthetic genes upregulated in *Atknat7* lines, while another shows downregulation (Li *et al.*, 2012; He *et al.*, 2018). AtKNAT7 protein can bind to the *AtIRX9* promoter, a gene responsible for xylan backbone synthesis.

The rice orthologue of *KNAT7*, *OsKNOR1* (also known as *OsKNAT7*), can negatively regulate cell wall synthesis in interfascicular fibre cells (Wang *et al.*, 2019; Zhao *et al.*, 2019). *Osknor1* mutants have thicker interfascicular fibre walls, with no reported xylem phenotype. However, *OsKNOR1* analysis revealed other functions unique to *AtKNAT7* (Wang *et al.*, 2019). OsKNOR1 protein interacts with OsSWN3 (also known as OsVND7 and OsNAC31) and OsGRF4 proteins and transient gene expression analysis showed that OsKNOR1–OsSWN3 jointly regulated *OsMYB61* and *OsMYB103* expression, with the addition of *OsKNOR1* reducing the positive regulation of OsSWN3 targets. Similarly, OsGRF4 is known to activate expression of expansin genes *OsEXPB3*, *OsEXPB17*, and *OsEXPA6* and addition of *OsKNOR1* also repressed that effect. This suggests that *OsKNOR1* regulates wall thickening and cell expansion by decreasing the transcriptional activation of *OsSWN3* and *OsGRF4*, respectively. This was validated by the observation of wall thickening in stem internodes and cell elongation along the panicle in relation to the expression of *OsKNOR1*, *OsSWN3* and *OsGRF4* (Wang *et al.*, 2019).

Among the genes co-expressed with *B. distachyon* *CSLF6*, a predominant MLG synthase, was a trihelix family transcription factor (*BdTHX1*) (Fan *et al.*, 2018; Kim *et al.*, 2018). This is the first THX protein associated with cell wall biosynthesis and the first shown to bind directly to a *CSLF* gene. *In planta* and *in vitro* assays showed that BdTHX1 protein binds to the GT element in the second intron of *BdCSLF6* and to the 3' region of glycoside hydrolase family 61 endotransglucosylase/hydrolase 8 (*BdXTH8*), a grass-specific

enzyme that uses MLG as a substrate (Fan *et al.*, 2018). Attempts to recover viable transgenic plants were unsuccessful and suggest a strong selection against the perturbation of *BdTHX1*, thus it is uncertain if it is a transcriptional activator or repressor.

The presence of phytohormone gibberellin results in the induction of secondary wall CESA genes in rice, and *OsMYB103* is necessary for that activation (Ye *et al.*, 2015). Similarly, the function of OsSWN2 (also known as OsNAC29) and OsSWN3 protein can be activated by gibberellins. The mechanism for activation is to degrade a protein interaction with the rice DELLA protein SLENDER RICE1 (SLR1) (Huang *et al.*, 2015). SLR1 protein is degraded in the presence of gibberellins and, subsequently, wall gene expression is activated (Fig. 5). A similar mechanism for gibberellin signalling in eudicots has not been reported and the role of this hormone in the regulation of wall thickening is not well resolved.

There is nearly complete overlap between the regulatory network components between eudicots and grasses. The distinctions between grass and eudicot walls are difficult to assign to differences in varying functions or members of the regulatory network. While no LBD, BLH or HD-ZIP III has been described as regulators of cell wall biosynthesis in grasses, it is possible that they have simply not been studied or reported. Additionally, *THX1* is likely to be unique to grasses as it regulates a hemicellulose gene not present in eudicots. Meta-analysis of microarray gene expression data, to make a combined mutual ranked network for rice and *A. thaliana*, has revealed differences in the relative importance of each regulator (Zhao *et al.*, 2019). The degree of connectivity among genes, which is the number of edges for each network node, can suggest the importance of each transcription factor. Some highly connected genes in *A. thaliana*, including *VND1/2/6/7* and *AtMYB46/83* have a two-fold to five-fold decrease in connectivity in rice. Conversely, transcription factors with considerably more connections in rice than *A. thaliana* are *OsSND2/3*, the rice orthologue of *KNAT7*, *KNOR1* as well as *OsSWN1*, the orthologue of *AtNST1*.

## XII. Conclusions

Much progress has been made in recent years to better understand grass cell wall composition and regulation, in large part thanks to the numerous genetic and genomic resources that have been developed. A case in point, *B. distachyon* as a model grass system has been central to these efforts, and provides fertile ground for future studies. The unique features of grass cell walls, such as MLG synthesis and the integration of hydroxycinnamates into lignin and xylan are beginning to be uncovered in detail. Elements that were thought to be more common between grasses and eudicots, such as lignin synthesis, continue to show evidence that there is yet unexplored diversity in plant cell wall chemistry, with alternate lignin biosynthetic pathways and atypical monomer components. Regulation remains an area of much overlap, but rather than playing catch-up with eudicots, grass networks now offer new insights that expand the cell wall network. Uncovering grass-specific functions, such as *BdTHX1* regulation of MLG, highlight the opportunities to advance this important area of plant biology.




## Acknowledgements

We apologise to our colleagues for not being able to cite all the relevant literature due to space limitations. This work was supported by The National Science Foundation Division of Integrative Organismal Systems (NSF IOS-1558072 to SPH).

## Author contributions

JHC, RS and SPH wrote the manuscript. RS and SPH contributed equally to this work.

## ORCID

Joshua H. Coomey  <https://orcid.org/0000-0001-5866-6671>  
 Samuel P. Hazen  <https://orcid.org/0000-0001-9131-8994>  
 Richard Sibout  <https://orcid.org/0000-0002-0639-5643>

## References

- Agarwal T, Grotewold E, Doseff AI, Gray J. 2016. MYB31/MYB42 syntelogs exhibit divergent regulation of phenylpropanoid genes in maize, sorghum and rice. *Scientific Reports* 6: 28502.
- Anders N, Wilkinson MD, Lovegrove A, Freeman J, Tryfona T, Pellny TK, Weimar T, Mortimer JC, Stott K, Baker JM *et al.* 2012. Glycosyl transferases in family 61 mediate arabinofuranosyl transfer onto xylan in grasses. *Proceedings of the National Academy of Sciences, USA* 109: 989–993.
- Arioli T, Peng L, Betzner AS, Burn J, Witte W, Herth W, Camilleri C, Hofte H, Plazinski J, Birch R *et al.* 1998. Molecular analysis of cellulose biosynthesis in *Arabidopsis*. *Science* 279: 717–720.
- Atmadojo MA, Hao Z, Mohnen D. 2013. Evolving views of pectin biosynthesis. *Annual Review of Plant Biology* 64: 747–779.
- Barriere Y, Ralph J, Mechin V, Guillaumie S, Grabber JH. 2004. Genetic and molecular basis of grass cell wall biosynthesis and degradability. II. Lessons from *brown-midrib* mutants. *Comptes Rendus Biologies* 327: 847.
- Barros J, Escamilla-Trevino L, Song L, Rao X, Serrani-Yarce JC, Palacios MD, Engle N, Choudhury FK, Tschaplinski TJ, Venables BJ *et al.* 2019. 4-Coumarate 3-hydroxylase in the lignin biosynthesis pathway is a cytosolic ascorbate peroxidase. *Nature Communications* 10: 1994.
- Barros J, Serrani-Yarce JC, Chen F, Baxter D, Venables BJ, Dixon RA. 2016. Role of bifunctional ammonia-lyase in grass cell wall biosynthesis. *Nature Plants* 2: 16050.
- Bartley LE, Peck ML, Kim S-R, Ebert B, Manisseri C, Chiniquy DM, Sykes R, Gao L, Rautengarten C, Vega-Sánchez ME *et al.* 2013. Overexpression of a BAH1 acyltransferase, *OsAt10*, alters rice cell wall hydroxycinnamic acid content and saccharification. *Plant Physiology* 161: 1615–1633.
- Bauer WD, Talmadge KW, Keegstra K, Albersheim P. 1973. The structure of plant cell walls: II. The hemicellulose of the walls of suspension-cultured sycamore cells. *Plant Physiology* 51: 174–187.
- Berthet S, Demont-Caulet N, Pollet B, Bidzinski P, Cézard L, Le Bris P, Borrega N, Hervé J, Blondet E, Balzergue S *et al.* 2011. Disruption of *LACCASE4* and 17 results in tissue-specific alterations to lignification of *Arabidopsis thaliana* stems. *Plant Cell* 23: 1124–1137.
- Betekhtin A, Milewska-Hendel A, Lusinska J, Chajec L, Kurczynska E, Hasterok R. 2018. Organ and tissue-specific localisation of selected cell wall epitopes in the zygotic embryo of *Brachypodium distachyon*. *International Journal of Molecular Sciences* 19: 275.
- Bhatia R, Dalton S, Roberts LA, Moron-Garcia OM, Iacono R, Kosik O, Gallagher JA, Bosch M. 2019. Modified expression of *ZmMYB167* in *Brachypodium distachyon* and *Zea mays* leads to increased cell wall lignin and phenolic content. *Scientific Reports* 9: 8800.
- Bogamuwa SP, Jang J-C. 2014. Tandem CCCH zinc finger proteins in plant growth, development and stress response. *Plant & Cell Physiology* 55: 1367–1375.

- Bokor B, Ondoš S, Vaculík M, Bokorová S, Weidinger M, Lichtscheidl I, Turňa J, Lux A. 2017. Expression of genes for Si uptake, accumulation and correlation of Si with other elements in ionome of maize kernel. *Frontiers in Plant Science* 8: 1063.
- Bouvier d'Yvoire M, Bouchabke-Coussa O, Voorend W, Antelme S, Cézard L, Légée F, Lebris P, Legay S, Whitehead C, McQueen-Mason SJ *et al.* 2013. Disrupting the *cinnamyl alcohol dehydrogenase 1 gene (BdCAD1)* leads to altered lignification and improved saccharification in *Brachypodium distachyon*. *The Plant Journal* 73: 496–508.
- Brabham C, Singh A, Stork J, Rong Y, Kumar I, Kikuchi K, Yingling YG, Brutnell TP, Rose JKC, Debolt S. 2019. Biochemical and physiological flexibility accompanies reduced cellulose biosynthesis in *Brachypodium cesa1<sub>S830N</sub>*. *Annals of Botany*. Plants 11: lz041.
- Bragg JN, Wu J, Gordon SP, Guttman ME, Thilmony R, Lazo GR, Gu YQ, Vogel JP. 2012. Generation and characterization of the Western Regional Research Center *Brachypodium* T-DNA insertional mutant collection. *PLoS ONE* 7: e41916.
- Bromley JR, Busse-Wicher M, Tryfona T, Mortimer JC, Zhang Z, Brown DM, Dupree P. 2013. GUX1 and GUX2 glucuronyltransferases decorate distinct domains of glucuronoxylan with different substitution patterns. *The Plant Journal* 74: 423–434.
- Brown DM, Zeef LAH, Ellis J, Goodacre R, Turner SR. 2005. Identification of novel genes in *Arabidopsis* involved in secondary cell wall formation using expression profiling and reverse genetics. *Plant Cell* 17: 2281–2295.
- Brown DM, Zhang ZN, Stephens E, Dupree P, Turner SR. 2009. Characterization of IRX10 and IRX10-like reveals an essential role in glucuronoxylan biosynthesis in *Arabidopsis*. *The Plant Journal* 57: 732.
- Buanafina MMO, Fescemyer HW, Sharma M, Shearer EA. 2016. Functional testing of a PF02458 homologue of putative rice arabinoxylan feruloyl transferase genes in *Brachypodium distachyon*. *Planta* 243: 659–674.
- Buliga GS, Brant DA, Fincher GB. 1986. The sequence statistics and solution conformation of a barley (1→3, 1→4)-beta-D-glucan. *Carbohydrate Research* 157: 139–156.
- Bulone V, Schwerdt JG, Fincher GB. 2019. Co-evolution of enzymes involved in plant cell wall metabolism in the grasses. *Frontiers in Plant Science* 10: 1009.
- Burton RA, Collins HM, Kibble NAJ, Smith JA, Shirley NJ, Jobling SA, Henderson M, Singh RR, Pettolino F, Wilson SM *et al.* 2011. Over-expression of specific *HvCslF* cellulose synthase-like genes in transgenic barley increases the levels of cell wall (1,3;1,4)-β-d-glucans and alters their fine structure. *Plant Biotechnology Journal* 9: 117–135.
- Burton RA, Fincher GB. 2012. Current challenges in cell wall biology in the cereals and grasses. *Frontiers in Plant Science* 3: 130.
- Burton RA, Fincher GB. 2014. Evolution and development of cell walls in cereal grains. *Frontiers in Plant Science* 5: 456.
- Burton RA, Gidley MJ, Fincher GB. 2010a. Heterogeneity in the chemistry, structure and function of plant cell walls. *Nature Chemical Biology* 6: 724–732.
- Burton RA, Ma G, Baumann U, Harvey AJ, Shirley NJ, Taylor J, Pettolino F, Bacic A, Beatty M, Simmons CR *et al.* 2010b. A customized gene expression microarray reveals that the brittle stem phenotype *fs2* of barley is attributable to a retroelement in the *HvCesA4* cellulose synthase gene. *Plant Physiology* 153: 1716–1728.
- Busse-Wicher M, Gomes TCF, Tryfona T, Nikolovski N, Stott K, Grantham NJ, Bolam DN, Skaf MS, Dupree P. 2014. The pattern of xylan acetylation suggests xylan may interact with cellulose microfibrils as a twofold helical screw in the secondary plant cell wall of *Arabidopsis thaliana*. *The Plant Journal* 79: 492–506.
- Carpita NC, McCann MC. 2010. The maize mixed-linkage (1→3), (1→4)-beta-D-glucan polysaccharide is synthesized at the golgi membrane. *Plant Physiology* 153: 1362–1371.
- Cass CL, Peraldi A, Dowd PF, Mottiar Y, Santoro N, Karlen SD, Bukhman YV, Foster CE, Thrower N, Bruno LC *et al.* 2015. Effects of *PHENYLALANINE AMMONIA LYASE (PAL)* knockdown on cell wall composition, biomass digestibility and biotic and abiotic stress responses in *Brachypodium*. *Journal of Experimental Botany* 66: 4317–4335.
- Cesarino I, Araújo P, Sampaio MR, Vicentini R, Berthet S, Demedts B, Vanholme B, Boerjan W, Mazzafera P. 2013. Expression of *SofLAC*, a new laccase in sugarcane, restores lignin content but not S:G ratio of *Arabidopsis lac17* mutant. *Journal of Experimental Botany* 64: 1769–1781.
- Chiniquy D, Sharma V, Schultink A, Baidoo EE, Rautengarten C, Cheng K, Carroll A, Ulvskov P, Harholt J, Keasling JD *et al.* 2012. XAX1 from glycosyltransferase family 61 mediates xylosyltransfer to rice xylan. *Proceedings of the National Academy of Sciences, USA* 109: 17117–17122.
- Christensen U, Alonso-Simon A, Scheller HV, Willats WG, Harholt J. 2010. Characterization of the primary cell walls of seedlings of *Brachypodium distachyon* - a potential model plant for temperate grasses. *Phytochemistry* 71: 62–69.
- Clarke HT, Gillespie HB, Weissshaus SZ. 1933. The action of formaldehyde on amines and amino acids. *Journal of the American Chemical Society* 55: 4571–4587.
- Coomey J, Hazen SP. 2015. *Brachypodium* as a model for the cell wall and biomass crops. In: Vogel JP, ed. *Plant genetics and genomics: crops and models. Genetics and genomics of Brachypodium*. New York, NY, USA: Springer-Verlag, 197–217.
- D'Auria JC. 2006. Acyltransferases in plants: a good time to be BAHD. *Current Opinion in Plant Biology* 9: 331–340.
- Dalmais M, Antelme S, Ho-Yue-Kuang S, Wang Y, Darracq O, Bouvier d'Yvoire M, Cézard L, Légée F, Blondet E, Oria N *et al.* 2013. A TILLING platform for functional genomics in *Brachypodium distachyon*. *PLoS ONE* 8: e65503.
- de Souza WR, Martins PK, Freeman J, Pellny TK, Michaelson LV, Sampaio BL, Vinecky F, Ribeiro AP, da Cunha BADB, Kobayashi AK *et al.* 2018. Suppression of a single BAHD gene in *Setaria viridis* causes large, stable decreases in cell wall feruloylation and increases biomass digestibility. *New Phytologist* 218: 81–93.
- de Souza WR, Pacheco TF, Duarte KE, Sampaio BL, de Oliveira Molinari PA, Martins PK, Santiago TR, Formighieri EF, Vinecky F, Ribeiro AP *et al.* 2019. Silencing of a BAHD acyltransferase in sugarcane increases biomass digestibility. *Biotechnology for Biofuels* 12: 111.
- del Río JC, Rencoret J, Gutiérrez A, Kim H, Ralph J. 2018. Structural characterization of lignin from maize (*Zea mays* L.) fibers: evidence for diferuloylputrescine incorporated into the lignin polymer in maize kernels. *Journal of Agricultural and Food Chemistry* 66: 4402–4413.
- del Río JC, Rencoret J, Prinsen P, Martínez ÁT, Ralph J, Gutiérrez A. 2012. Structural characterization of wheat straw lignin as revealed by analytical pyrolysis, 2D-NMR and reductive cleavage methods. *Journal of Agricultural and Food Chemistry* 60: 5922–5935.
- Deshmukh RK, Ma JF, Bélanger RR. 2017. Editorial: Role of silicon in plants. *Frontiers in Plant Science* 8: 1858.
- Dimitroff G, Little A, Lahnstein J, Schwerdt JG, Srivastava V, Bulone V, Burton RA, Fincher GB. 2016. (1,3;1,4)-β-glucan biosynthesis by the CSLF6 enzyme: position and flexibility of catalytic residues influence product fine structure. *Biochemistry* 55: 2054–2061.
- Do C-T, Pollet B, Thévenin J, Sibout R, Denoue D, Barrière Y, Lapiere C, Jouanin L. 2007. Both caffeoyl Coenzyme A 3-O-methyltransferase 1 and caffeic acid O-methyltransferase 1 are involved in redundant functions for lignin, flavonoids and sinapoyl malate biosynthesis in *Arabidopsis*. *Planta* 226: 1117–1129.
- Eloy NB, Voorend W, Lan W, Saleme MLS, Cesarino I, Vanholme R, Smith RA, Goeminne G, Pallidis A, Morreel K *et al.* 2017. Silencing *CHALCONE SYNTHASE* in maize impedes the incorporation of tricetin into lignin and increases lignin content. *Plant Physiology* 173: 998–1016.
- Esau K. 1977. *Anatomy of seed plants*. New York, NY, USA: John Wiley & Sons.
- Eudes A, Dutta T, Deng K, Jacquet N, Sinha A, Benites VT, Baidoo EEK, Richel A, Sattler SE, Northen TR *et al.* 2017. SbcOMT (Bmr12) is involved in the biosynthesis of tricetin-lignin in sorghum. *PLoS ONE* 12: e0178160.
- Fan M, Herburger K, Jensen JK, Zemelis-Durfee S, Brandizzi F, Fry SC, Wilkerson CG. 2018. A trihelix family transcription factor is associated with key genes in mixed-linkage glucan accumulation. *Plant Physiology* 178: 1207–1221.
- Fincher GB. 2009. Exploring the evolution of (1,3;1,4)-beta-D-glucans in plant cell walls: comparative genomics can help!. *Current Opinion in Plant Biology* 12: 140–147.
- Floyd SK, Bowman JL. 2006. Distinct developmental mechanisms reflect the independent origins of leaves in vascular plants. *Current Biology* 16: 1911–1917.
- Fornalé S, Rencoret J, García-Calvo L, Encina A, Rigau J, Gutiérrez A, Del Río JC, Caparros-Ruiz D. 2017. Changes in cell wall polymers and degradability in maize mutants lacking 3'- and 5'-O-methyltransferases involved in lignin biosynthesis. *Plant & Cell Physiology* 58: 240–255.
- Fornalé S, Shi X, Chai C, Encina A, Irar S, Capellades M, Fuguet E, Torres J-L, Rovira P, Puigdomènech P *et al.* 2010. ZmMYB31 directly represses maize lignin genes and redirects the phenylpropanoid metabolic flux. *The Plant Journal* 64: 633–644.



- Fornalé S, Sonbol F-M, Maes T, Capellades M, Puigdomènech P, Rigau J, Caparrós-Ruiz D. 2006. Down-regulation of the maize and *Arabidopsis thaliana* caffeic acid *O*-methyl-transferase genes by two new maize R2R3-MYB transcription factors. *Plant Molecular Biology* 62: 809–823.
- Francin-Allami M, Alvarado C, Daniel S, Geairon A, Saulnier L, Guillon F. 2019. Spatial and temporal distribution of cell wall polysaccharides during grain development of *Brachypodium distachyon*. *Plant Science* 280: 367–382.
- Fry SC. 1989. The structure and functions of xyloglucan. *Journal of Experimental Botany* 40: 1–11.
- Gibeaut DM, Pauly M, Bacic A, Fincher GB. 2005. Changes in cell wall polysaccharides in developing barley (*Hordeum vulgare*) coleoptiles. *Planta* 221: 729–738.
- Głazowska S, Baldwin L, Mravec J, Bukh C, Hansen TH, Jensen MM, Fangel JU, Willats WGT, Glasius M, Felby C *et al.* 2018a. The impact of silicon on cell wall composition and enzymatic saccharification of *Brachypodium distachyon*. *Biotechnology for Biofuels* 11: 171.
- Głazowska S, Murozuka E, Persson DP, Castro PH, Schjoerring JK. 2018b. Silicon affects seed development and leaf macrohair formation in *Brachypodium distachyon*. *Physiologia Plantarum* 163: 231–246.
- Gordon SP, Contreras-Moreira B, Woods DP, Des Marais DL, Burgess D, Shu S, Stritt C, Roulin AC, Schackwitz W, Tyler L *et al.* 2017. Extensive gene content variation in the *Brachypodium distachyon* pan-genome correlates with population structure. *Nature Communications* 8: 2184.
- Granier F, Lemaire A, Wang Y, LeBris P, Antelme S, Vogel J, Laudencia-Chinguanco D, Sibout R. 2015. Chemical and radiation mutagenesis: induction and detection by whole genome sequencing. In: Vogel JP, ed. *Plant genetics and genomics: crops and models. Genetics and genomics of Brachypodium*. New York, NY, USA: Springer-Verlag, 155–170.
- Guillon F, Bouchet B, Jamme F, Robert P, Quémener B, Barron C, Larré C, Dumas P, Saulnier L. 2011. *Brachypodium distachyon* grain: characterization of endosperm cell walls. *Journal of Experimental Botany* 62: 1001–1015.
- Ha CM, Escamilla-Trevino L, Yarce JCS, Kim H, Ralph J, Chen F, Dixon RA. 2016. An essential role of caffeoyl shikimate esterase in monolignol biosynthesis in *Medicago truncatula*. *The Plant Journal* 86: 363–375.
- Handakumbura PP, Brow K, Whitney IP, Zhao K, Sanguinet KA, Lee SJ, Olins J, Romero-Gamboá SP, Harrington MJ, Bascom CJ *et al.* 2018. SECONDARY WALL ASSOCIATED MYB1 is a positive regulator of secondary cell wall thickening in *Brachypodium distachyon* and is not found in the Brassicaceae. *The Plant Journal* 96: 485–699.
- Handakumbura P, Matos D, Osmont K, Harrington M, Heo K, Kafle K, Kim S, Baskin T, Hazen S. 2013. Perturbation of *Brachypodium distachyon* CELLULOSE SYNTHASE A4 or 7 results in abnormal cell walls. *BMC Plant Biology* 13: 131.
- Hatfield RD, Marita JM, Frost K, Grabber J, Ralph J, Lu F, Kim H. 2009. Grass lignin acylation: *p*-coumaroyl transferase activity and cell wall characteristics of C<sub>3</sub> and C<sub>4</sub> grasses. *Planta* 229: 1253–1267.
- Hatfield RD, Rancour DM, Marita JM. 2016. Grass cell walls: a story of cross-linking. *Frontiers in Plant Science* 7: 2056.
- Hattori T, Inanaga S, Araki H, An P. 2005. Application of silicon enhanced drought tolerance in *Sorghum bicolor*. *Physiologia Plantarum* 123: 359–474.
- He J-B, Zhao X-H, Du P-Z, Zeng W, Beahan CT, Wang Y-Q, Li H-L, Bacic A, Wu A-M. 2018. KNAT7 positively regulates xylan biosynthesis by directly activating IRX9 expression in *Arabidopsis*. *Journal of Integrative Plant Biology* 60: 514–528.
- Herbaut M, Zoghliani A, Habrant A, Falourd X, Foucat L, Chabbert B, Paës G. 2018. Multimodal analysis of pretreated biomass species highlights generic markers of lignocellulose recalcitrance. *Biotechnology for Biofuels* 11: 52.
- Higuchi T, Ito Y, Kawamura I. 1967. *p*-hydroxyphenylpropane component of grass lignin and role of tyrosine-ammonia lyase in its formation. *Phytochemistry* 6: 875–881.
- Hirano K, Kondo M, Aya K, Miyao A, Sato Y, Antonio BA, Namiki N, Nagamura Y, Matsuoka M. 2013. Identification of transcription factors involved in rice secondary cell wall formation. *Plant & Cell Physiology* 54: 1791–1802.
- Høj PB, Fincher GB. 1995. Molecular evolution of plant beta-glucan endohydrolases. *The Plant Journal* 7: 367–379.
- Huang D, Wang S, Zhang B, Shang-Guan K, Shi Y, Zhang D, Liu X, Wu K, Xu Z, Fu X *et al.* 2015. A gibberellin-mediated DELLA-NAC signaling cascade regulates cellulose synthesis in rice. *Plant Cell* 27: 1681–1696.
- Huang P, Yoshida H, Yano K, Kinoshita S, Kawai K, Koketsu E, Hattori M, Takehara S, Huang J, Hirano K *et al.* 2018. OsIDD2, a zinc finger and INDETERMINATE DOMAIN protein, regulates secondary cell wall formation. *Journal of Integrative Plant Biology* 60: 130–143.
- Jensen JK, Wilkerson CG. 2017. *Brachypodium* as an experimental system for the study of stem parenchyma biology in grasses. *PLoS ONE* 12: e0173095.
- Jobling SA. 2015. Membrane pore architecture of the CslF6 protein controls (1–3,1–4)- $\beta$ -glucan structure. *Science Advances* 1: e1500069.
- Kapp N, Barnes WJ, Richard TL, Anderson CT. 2015. Imaging with the fluorogenic dye Basic Fuchsin reveals subcellular patterning and ecotype variation of lignification in *Brachypodium distachyon*. *Journal of Experimental Botany* 66: 4295–4304.
- Karlen SD, Zhang C, Peck ML, Smith RA, Padmakshan D, Helmich KE, Free HCA, Lee S, Smith BG, Lu F *et al.* 2016. Monolignol ferulate conjugates are naturally incorporated into plant lignins. *Science Advances* 2: e1600393.
- Kido N, Yokoyama R, Yamamoto T, Furukawa J, Iwai H, Satoh S, Nishitani K. 2015. The matrix polysaccharide (1;3,1;4)- $\beta$ -D-glucan is involved in silicon-dependent strengthening of rice cell wall. *Plant & Cell Physiology* 56: 1679.
- Kim S-J, Zemelis S, Keegstra K, Brandizzi F. 2015. The cytoplasmic localization of the catalytic site of CSLF6 supports a channeling model for the biosynthesis of mixed-linkage glucan. *The Plant Journal* 81: 537–547.
- Kim S-J, Zemelis-Durfee S, Jensen JK, Wilkerson CG, Keegstra K, Brandizzi F. 2018. In the grass species *Brachypodium distachyon*, the production of mixed-linkage (1,3; 1,4)- $\beta$ -glucan (MLG) occurs in the Golgi apparatus. *The Plant Journal* 93: 1062–1075.
- Kim W-C, Ko J-H, Han K-H. 2012. Identification of a *cis*-acting regulatory motif recognized by MYB46, a master transcriptional regulator of secondary wall biosynthesis. *Plant Molecular Biology* 78: 489–501.
- Ko J-H, Kim W-C, Han K-H. 2009. Ectopic expression of MYB46 identifies transcriptional regulatory genes involved in secondary wall biosynthesis in *Arabidopsis*. *The Plant Journal* 60: 649–665.
- Kotake T, Aohara T, Hirano K, Sato A, Kaneko Y, Tsumuraya Y, Takatsuji H, Kawasaki S. 2011. Rice *Brittle culm 6* encodes a dominant-negative form of CesaA protein that perturbs cellulose synthesis in secondary cell walls. *Journal of Experimental Botany* 62: 2053–2062.
- Kubo M, Udagawa M, Nishikubo N, Horiguchi G, Yamaguchi M, Ito J, Mimura T, Fukuda H, Demura T. 2005. Transcription switches for protoxylem and metaxylem vessel formation. *Genes & Development* 19: 1855–1860.
- Lam PY, Tobimatsu Y, Matsumoto N, Suzuki S, Lan W, Takeda Y, Yamamura M, Sakamoto M, Ralph J, Lo C *et al.* 2019. OsCALD1 is a bifunctional *O*-methyltransferase involved in the biosynthesis of tricin-lignins in rice cell walls. *Scientific Reports* 9: 11597.
- Lam PY, Tobimatsu Y, Takeda Y, Suzuki S, Yamamura M, Umezawa T, Lo C. 2017. Disrupting flavone synthase II alters lignin and improves biomass digestibility. *Plant Physiology* 174: 972–985.
- Lan W, Lu F, Regner M, Zhu Y, Rencoret J, Ralph SA, Zakai UI, Morreel K, Boerjan W, Ralph J. 2015. Tricin, a flavonoid monomer in monocot lignification. *Plant Physiology* 167: 1284–1295.
- Lan W, Rencoret J, Lu F, Karlen SD, Smith BG, Harris PJ, Del Río JC, Ralph J. 2016. Tricin-lignins: occurrence and quantitation of tricin in relation to phylogeny. *The Plant Journal* 88: 1046–1057.
- Lan W, Yue F, Rencoret J, Del Río JC, Boerjan W, Lu F, Ralph J. 2018. Elucidating tricin-lignin structures: assigning correlations in HSQC spectra of monocot lignins. *Polymers* 10: 916.
- Langer RHM. 1979. *How grasses grow*. London, UK: Edward Arnold.
- Lapierre C, Voxel A, Boutet S, Ralph J. 2019. Arabinose conjugates diagnostic of ferulate-ferulate and ferulate-monolignol cross-coupling are released by mild acidolysis of grass cell walls. *Journal of Agricultural and Food Chemistry* 67: 12962–12971.
- Lazaridou A, Biliaderis CG. 2007. Molecular aspects of cereal  $\beta$ -glucan functionality: physical properties, technological applications and physiological effects. *Journal of Cereal Science* 46: 101–118.
- Le Bris P, Wang Y, Barbereau C, Antelme S, Cézard L, Legée F, D'Orlando A, Dalmais M, Bendahmane A, Schuetz M *et al.* 2019. Inactivation of LACCASE8 and LACCASE5 genes in *Brachypodium distachyon* leads to severe decrease in lignin content and high increase in saccharification yield without impacting plant integrity. *Biotechnology for Biofuels* 12: 181.



- Lee C, Teng Q, Zhong R, Yuan Y, Ye Z-H. 2014. Functional roles of rice glycosyltransferase family GT43 in xylan biosynthesis. *Plant Signaling & Behavior* 9: e27809.
- Lee CH, O'Neill MA, Tsumuraya Y, Darvill AG, Ye ZH. 2007. The *irregular xylem9* mutant is deficient in xylan xylosyltransferase activity. *Plant & Cell Physiology* 48: 1624.
- Lee K-H, Du Q, Zhuo C, Qi L, Wang H. 2019. LBD29-involved auxin signaling represses NAC master regulators and fiber wall biosynthesis. *Plant Physiology* 181: 595–608.
- Lee SJ, Warnick TA, Pattathil S, Alvelo-Maurosa JG, Serapiglia MJ, McCormick H, Brown V, Young NF, Schnell DJ, Smart LB *et al.* 2012. Biological conversion assay using *Clostridium phytofermentans* to estimate plant feedstock quality. *Biotechnology for biofuels* 5: 5.
- Li E, Bhargava A, Qiang W, Friedmann MC, Forneris N, Savidge RA, Johnson LA, Mansfield SD, Ellis BE, Douglas CJ. 2012. The Class II *KNOX* gene *KNAT7* negatively regulates secondary wall formation in *Arabidopsis* and is functionally conserved in *Populus*. *New Phytologist* 194: 102–115.
- Little A, Schwerdt JG, Shirley NJ, Khor SF, Neumann K, O'Donovan LA, Lahnstein J, Collins HM, Henderson M, Fincher GB *et al.* 2018. Revised phylogeny of the cellulose synthase gene superfamily: Insights into cell wall evolution. *Plant Physiology* 177: 1124–1141.
- Liu D, Zehfroosh N, Hancock BL, Hines K, Fang W, Kilfoil M, Learned-Miller E, Sanguinet KA, Goldner LS, Baskin TI. 2017. Imaging cellulose synthase motility during primary cell wall synthesis in the grass *Brachypodium distachyon*. *Scientific Reports* 7: 15111.
- Liu L, Hsia MM, Dama M, Vogel J, Pauly M. 2016. A xyloglucan backbone 6-O-acetyltransferase from *Brachypodium distachyon* modulates xyloglucan xylosylation. *Molecular Plant* 9: 615–617.
- Ma JF, Yamaji N. 2006. Silicon uptake and accumulation in higher plants. *Trends in Plant Science* 11: 392–397.
- Ma JF, Yamaji N, Tamai K, Mitani N. 2007. Genotypic difference in silicon uptake and expression of silicon transporter genes in rice. *Plant Physiology* 145: 919–924.
- MacKinnon KJ-M, Cole BJ, Yu C, Coomey JH, Hartwick NT, Remigereau M-S, Duffy T, Michael TP, Kay SA, Hazen SP. 2020. Changes in ambient temperature are the prevailing cue in determining *Brachypodium distachyon* diurnal gene regulation. *New Phytologist*. doi: 10.1111/nph.16507
- Marita JM, Hatfield RD, Rancour DM, Frost KE. 2014. Identification and suppression of the *p*-coumaroyl CoA:hydroxycinnamyl alcohol transferase in *Zea mays* L. *The Plant Journal* 78: 850–864.
- Mariotti PE, Sibout R, Lapierre C, Fangel JU, Willats WGT, Hofte H, Gómez LD, McQueen-Mason SJ. 2014. Range of cell-wall alterations enhance saccharification in *Brachypodium distachyon* mutants. *Proceedings of the National Academy of Sciences, India. Section A. Physical Sciences* 111: 14601–14606.
- Martínez-Abad A, Berglund J, Toriz G, Gatenholm P, Henriksson G, Lindström M, Wohler J, Vilaplana F. 2017. Regular motifs in xylan modulate molecular flexibility and interactions with cellulose surfaces. *Plant Physiology* 175: 1579–1592.
- Matos DA, Whitney IP, Harrington MJ, Hazen SP. 2013. Cell walls and the developmental anatomy of the *Brachypodium distachyon* stem internode. *PLoS ONE* 8: e80640.
- Méchin V, Laluc A, Legée F, Cézard L, Denoue D, Barrière Y, Lapierre C. 2014. Impact of the brown-midrib *bm5* mutation on maize lignins. *Journal of Agricultural and Food Chemistry* 62: 5102–5107.
- Mitani-Ueno N, Yamaji N, Ma JF. 2016. High silicon accumulation in the shoot is required for down-regulating the expression of Si transporter genes in rice. *Plant & Cell Physiology* 57: 2510–2518.
- Mitchell RAC, Dupree P, Shewry PR. 2007. A novel bioinformatics approach identifies candidate genes for the synthesis and feruloylation of arabinoxylan. *Plant Physiology* 144: 43–53.
- Mitsuda N, Iwase A, Yamamoto H, Yoshida M, Seki M, Shinozaki K, Ohme-Takagi M. 2007. NAC transcription factors, NST1 and NST3, are key regulators of the formation of secondary walls in woody tissues of *Arabidopsis*. *Plant Cell* 19: 270–280.
- Miyamoto T, Takada R, Tobimatsu Y, Takeda Y, Suzuki S, Yamamura M, Osakabe K, Osakabe Y, Sakamoto M, Umezawa T. 2019. OsMYB108 loss-of-function enriches *p*-coumaroylated and tricin lignin units in rice cell walls. *The Plant Journal* 98: 975–987.
- Mnich E, Bjarnholt N, Eudes A, Harholt J, Holland C, Jørgensen B, Larsen FH, Liu M, Manat R, Meyer AS *et al.* 2020. Phenolic cross-links: building and deconstructing the plant cell wall. *Natural Product Reports*. doi: 10.1039/c9np00028c
- Nakano Y, Yamaguchi M, Endo H, Rejab NA, Ohtani M. 2015. NAC-MYB-based transcriptional regulation of secondary cell wall biosynthesis in land plants. *Frontiers in Plant Science* 6: 288.
- Nemeth C, Freeman J, Jones HD, Sparks C, Pellny TK, Wilkinson MD, Dunwell J, Andersson AAM, Aman P, Guillon F *et al.* 2010. Down-regulation of the *CSLF6* gene results in decreased (1,3;1,4)-[beta]-D-glucan in endosperm of wheat. *Plant Physiology* 152: 1209–1218.
- Noda S, Koshiba T, Hattori T, Yamaguchi M, Suzuki S, Umezawa T. 2015. The expression of a rice secondary wall-specific cellulose synthase gene, *OsCesA7*, is directly regulated by a rice transcription factor, OsMYB58/63. *Planta* 242: 589–600.
- O'Malley RC, Huang S-SC, Song L, Lewsey MG, Bartlett A, Nery JR, Galli M, Gallavotti A, Ecker JR. 2016. Cistrome and epicistrome features shape the regulatory DNA landscape. *Cell* 165: 1280–1292.
- Ohashi-Ito K, Iwamoto K, Fukuda H. 2018. LOB DOMAIN-CONTAINING PROTEIN 15 positively regulates expression of *VND7*, a master regulator of tracheary elements. *Plant & Cell Physiology* 59: 989–996.
- Ohtani M, Nishikubo N, Xu B, Yamaguchi M, Mitsuda N, Goué N, Shi F, Ohme-Takagi M, Demura T. 2011. A NAC domain protein family contributing to the regulation of wood formation in poplar. *The Plant Journal* 67: 499–512.
- Olins JR, Lin L, Lee SJ, Trabucco GM, MacKinnon KJM, Hazen SP. 2018. Secondary wall regulating NACs differentially bind at the promoter at a *CELLULOSE SYNTHASE A4* cis-eQTL. *Frontiers in Plant Science* 9: 1895.
- Opanowicz M, Hands P, Betts D, Parker ML, Toole GA, Mills ENC, Doonan JH, Drea S. 2011. Endosperm development in *Brachypodium distachyon*. *Journal of Experimental Botany* 62: 735–748.
- Paredes A, Somerville CR, Ehrhardt D. 2006. Dynamic visualization of cellulose synthase demonstrates functional association with cortical microtubules. *Science* 312: 1491.
- Pear JR, Kawagoe Y, Schreckengost WE, Delmer DP, Stalker DM. 1996. Higher plants contain homologs of the bacterial celA genes encoding the catalytic subunit of cellulose synthase. *Proceedings of the National Academy of Sciences, USA* 93: 12637.
- Peña MJ, Zhong R, Zhou G-K, Richardson EA, O'Neill MA, Darvill AG, York WS, Ye Z-H. 2007. *Arabidopsis irregular xylem8* and *irregular xylem9*: implications for the complexity of glucuronoxylan biosynthesis. *Plant Cell* 19: 549–563.
- Perkins M, Smith RA, Samuels L. 2019. The transport of monomers during lignification in plants: anything goes but how? *Current Opinion in Biotechnology* 56: 69–74.
- Perry CC, Lu Y. 1992. Preparation of silicas from silicon complexes: role of cellulose in polymerisation and aggregation control. *Journal of the Chemical Society, Faraday Transactions* 88: 2915–2921.
- Persson S, Paredes A, Carroll A, Palsdottir H, Doblin M, Poindexter P, Khitrov N, Auer M, Somerville CR. 2007. Genetic evidence for three unique components in primary cell-wall cellulose synthase complexes in *Arabidopsis*. *Proceedings of the National Academy of Sciences, USA* 104: 15566–15571.
- Persson S, Wei H, Milne J, Page GP, Somerville CR. 2005. Identification of genes required for cellulose synthesis by regression analysis of public microarray data sets. *Proceedings of the National Academy of Sciences, USA* 102: 8633–8638.
- Petrik DL, Cass CL, Padmakshan D, Foster CE, Vogel JP, Karlen SD, Ralph J, Sedbrook JC. 2016. *BdCESA7*, *BdCESA8* and *BdPMT* utility promoter constructs for targeted expression to secondary cell-wall-forming cells of grasses. *Frontiers in Plant Science* 7: 55.
- Petrik DL, Karlen SD, Cass CL, Padmakshan D, Lu F, Liu S, Le Bris P, Antelme S, Santoro N, Wilkerson CG *et al.* 2014. *p*-Coumaroyl-CoA:monoglucuronol transferase (PMT) acts specifically in the lignin biosynthetic pathway in *Brachypodium distachyon*. *The Plant Journal* 77: 713–726.
- Pillonel C, Mulder MM, Boon JJ, Forster B, Binder A. 1991. Involvement of cinnamyl-alcohol dehydrogenase in the control of lignin formation in *Sorghum bicolor* L. Moench. *Planta* 185: 538–544.

- Piquemal J, Chamayou S, Nadaud I, Beckert M, Barriere Y, Mila I, Lapierre C, Rigau J, Puigdomenech P, Jauneau A *et al.* 2002. Down-regulation of caffeic acid O-methyltransferase in maize revisited using a transgenic approach. *Plant Physiology* 130: 1675–1685.
- Piston F, Uauy C, Fu L, Langston J, Labavitch J, Dubcovsky J. 2010. Down-regulation of four putative arabinoxylan feruloyl transferase genes from family PF02458 reduces ester-linked ferulate content in rice cell walls. *Planta* 231: 677–691.
- Polko JK, Kieber JJ. 2019. The regulation of cellulose biosynthesis in plants. *Plant Cell* 31: 282–296.
- Pyo H, Demura T, Fukuda H. 2007. TERE; a novel cis-element responsible for a coordinated expression of genes related to programmed cell death and secondary wall formation during differentiation of tracheary elements. *The Plant Journal* 51: 955–965.
- Ralph J, Grabber JH, Hatfield RD. 1995. Lignin-ferulate cross-links in grasses: active incorporation of ferulate polysaccharide esters into ryegrass lignins. *Carbohydrate Research* 275: 167–178.
- Ralph J, Hatfield RD, Grabber JH, Jung H-JG, Quideau S, Helm RF. 1998. Cell wall cross-linking in grasses by ferulates and diferulates. In: Lewis NG, Sarkanen S, eds. *ACS Symposium Series. Lignin and lignan biosynthesis*. Washington, DC, USA: American Chemical Society, 209–236.
- Ralph J, Quideau S, Grabber JH, Hatfield RD. 1994. Identification and synthesis of new ferulic acid dehydromers present in grass cell walls. *Journal of the Chemical Society*. 3485–3498.
- Rancour D, Marita J, Hatfield RD. 2012. Cell wall composition throughout development for the model grass *Brachypodium distachyon*. *Frontiers in Plant Science* 3: 266.
- Rao X, Chen X, Shen H, Ma Q, Li G, Tang Y, Pena M, York W, Frazier TP, Lenaghan S *et al.* 2019. Gene regulatory networks for lignin biosynthesis in switchgrass (*Panicum virgatum*). *Plant Biotechnology Journal* 17: 580–593.
- Rao X, Dixon RA. 2018. Current models for transcriptional regulation of secondary cell wall biosynthesis in grasses. *Frontiers in Plant Science* 9: 399.
- Roulin S, Buchala AJ, Fincher GB. 2002. Induction of (1→3,1→4)-beta-D-glucan hydrolases in leaves of dark-incubated barley seedlings. *Planta* 215: 51–59.
- Scheller HV, Ulvskov P. 2010. Hemicelluloses. *Annual Review of Plant Biology* 61: 263–289.
- Scholthof K-BG, Irigoyen S, Catalan P, Mandadi KK. 2018. *Brachypodium*: a monocot grass model genus for plant biology. *Plant Cell* 30: 1673.
- Scully ED, Gries T, Sarath G, Palmer NA, Baird L, Serapiglia MJ, Dien BS, Boateng AA, Ge Z, Funnell-Harris DL *et al.* 2016. Overexpression of *SbMyb60* impacts phenylpropanoid biosynthesis and alters secondary cell wall composition in *Sorghum bicolor*. *The Plant Journal* 85: 378–395.
- Shen H, He X, Poovaiya CR, Wuddineh WA, Ma J, Mann DGJ, Wang H, Jackson L, Tang Y, Neal Stewart C *et al.* 2012. Functional characterization of the switchgrass (*Panicum virgatum*) R2R3-MYB transcription factor *PvMYB4* for improvement of lignocellulosic feedstocks. *New Phytologist* 193: 121–136.
- Sibout R, Le Bris P, Legée F, Cézard L, Renault H, Lapierre C. 2016. Structural redesigning Arabidopsis lignins into alkali-soluble lignins through the expression of *p-coumaroyl-CoA:monolignol transferase* PMT. *Plant Physiology* 170: 1358–1366.
- Sibout R, Proost S, Hansen BO, Vaid N, Giorgi FM, Ho-Yue-Kuang S, Legée F, Cézard L, Bouchabké-Coussa O, Soulhat C *et al.* 2017. Expression atlas and comparative coexpression network analyses reveal important genes involved in the formation of lignified cell wall in *Brachypodium distachyon*. *New Phytologist* 215: 1009–1025.
- Simmons TJ, Mortimer JC, Bernardinelli OD, Pöppler A-C, Brown SP, Deazevedo ER, Dupree R, Dupree P. 2016. Folding of xylan onto cellulose fibrils in plant cell walls revealed by solid-state NMR. *Nature Communications* 7: 13902.
- Sindhu A, Langewisch T, Olek A, Multani DS, McCann MC, Vermerris W, Carpita NC, Johal G. 2007. Maize *brittle stalk2* encodes a COBRA-like protein expressed in early organ development but required for tissue flexibility at maturity. *Plant Physiology* 145: 1444–1459.
- Sonbol F-M, Fornalé S, Capellades M, Encina A, Touriño S, Torres J-L, Rovira P, Ruel K, Puigdomenech P, Rigau J *et al.* 2009. The maize *ZmMYB42* represses the phenylpropanoid pathway and affects the cell wall structure, composition and degradability in *Arabidopsis thaliana*. *Plant Molecular Biology* 70: 283–296.
- Soyano T, Thitamadee S, Machida Y, Chua N-H. 2008. *ASYMMETRIC LEAVES2-LIKE19/LATERAL ORGAN BOUNDARIES DOMAIN30* and *ASL20/LBD18* regulate tracheary element differentiation in *Arabidopsis*. *Plant Cell* 20: 3359–3373.
- Stewart JJ, Akiyama T, Chapple C, Ralph J, Mansfield SD. 2009. The effects on lignin structure of overexpression of ferulate 5-hydroxylase in hybrid poplar. *Plant physiology* 150: 621–635.
- Takeda Y, Tobimatsu Y, Karlen SD, Koshiba T, Suzuki S, Yamamura M, Murakami S, Mukai M, Hattori T, Osakabe K *et al.* 2018. Downregulation of *p-COUMAROYL ESTER 3-HYDROXYLASE* in rice leads to altered cell wall structures and improves biomass saccharification. *The Plant Journal* 95: 796–811.
- Tamasloukht B, Wong Quai Lam MS-J, Martinez Y, Tozo K, Barbier O, Jourda C, Jauneau A, Borderies G, Balzergue S, Renou J-P *et al.* 2011. Characterization of a cinnamoyl-CoA reductase 1 (CCR1) mutant in maize: effects on lignification, fibre development and global gene expression. *Journal of Experimental Botany* 62: 3837–3848.
- Tanaka K, Murata K, Yamazaki M, Onosato K, Miyao A, Hirochika H. 2003. Three distinct rice cellulose synthase catalytic subunit genes required for cellulose synthesis in the secondary wall. *Plant Physiology* 133: 73–83.
- Taylor-Teeple M, Lin L, de Lucas M, Turco G, Toal TW, Gaudinier A, Young NE, Trabucco GM, Veling MT, Lamothe R *et al.* 2015. An *Arabidopsis* gene regulatory network for secondary cell wall synthesis. *Nature* 517: 571–575.
- Trabucco GM, Matos DA, Lee SJ, Saathoff AJ, Priest HD, Mockler TC, Sarath G, Hazen SP. 2013. Functional characterization of cinnamyl alcohol dehydrogenase and caffeic acid O-methyltransferase in *Brachypodium distachyon*. *BMC Biotechnology* 13: 61.
- Trafford K, Haleux P, Henderson M, Parker M, Shirley NJ, Tucker MR, Fincher GB, Burton RA. 2013. Grain development in *Brachypodium* and other grasses: possible interactions between cell expansion, starch deposition and cell-wall synthesis. *Journal of Experimental Botany* 64: 5033–5047.
- Trethewey JAK, Campbell LM, Harris PJ. 2005. (1→3), (1→4)-β-D-glucans in the cell walls of the *Poales* (*sensu lato*): an immunogold labeling study using a monoclonal antibody. *American Journal of Botany* 92: 1660–1674.
- Trethewey JAK, Harris PJ. 2002. Location of (1→3)- and (1→3), (1→4)-beta-D-glucans in vegetative cell walls of barley (*Hordeum vulgare*) using immunogold labelling. *New Phytologist* 154: 347–358.
- Tyler L, Fangel JU, Fagerström AD, Steinwand MA, Raab TK, Willats WG, Vogel JP. 2014. Selection and phenotypic characterization of a core collection of *Brachypodium distachyon* inbred lines. *BMC Plant Biology* 14: 25.
- Urbanowicz BR, Peña MJ, Ratnaparkhe S, Avci U, Backe J, Steet HF, Foston M, Li H, O'Neill MA, Ragauskas AJ *et al.* 2012. 4-O-methylation of glucuronic acid in *Arabidopsis* glucuronoxylan is catalyzed by a domain of unknown function family 579 protein. *Proceedings of the National Academy of Sciences, USA* 109: 14253–14258.
- Valdivia ER, Herrera MT, Gianzo C, Fidalgo J, Revilla G, Zarra I, Sampedro J. 2013. Regulation of secondary wall synthesis and cell death by NAC transcription factors in the monocot *Brachypodium distachyon*. *Journal of Experimental Botany* 64: 1333–1343.
- Vanholme R, De Meester B, Ralph J, Boerjan W. 2019. Lignin biosynthesis and its integration into metabolism. *Current Opinion in Biotechnology* 56: 230–239.
- Vanholme R, Storme V, Vanholme B, Sundin L, Christensen JH, Goeminne G, Halpin C, Rohde A, Morreel K, Boerjan W. 2012. A systems biology view of responses to lignin biosynthesis perturbations in *Arabidopsis*. *Plant Cell* 24: 3506–3529.
- Varghese JN, Garrett TP, Colman PM, Chen L, Høj PB, Fincher GB. 1994. Three-dimensional structure of two plant beta-glucan endohydrolases with distinct substrate specificities. *Proceedings of the National Academy of Sciences, USA* 91: 2785–2789.
- Vega-Sánchez ME, Loqué D, Lao J, Catena M, Verhertbruggen Y, Herter T, Yang F, Harholt J, Ebert B, Baidoo EEK *et al.* 2015. Engineering temporal accumulation of a low recalcitrance polysaccharide leads to increased C6 sugar content in plant cell walls. *Plant Biotechnology Journal* 13: 903–914.
- Vega-Sanchez ME, Verhertbruggen Y, Christensen J, Chen X, Sharma V, Varanasi P, Jobling SA, Talbot M, White RG, Joo M *et al.* 2012. Loss of *Cellulose Synthase-like F6* function affects mixed-linkage glucan deposition, cell wall mechanical properties and defense responses in vegetative tissues of rice. *Plant Physiology* 159: 56–69.

- Vélez-Bermúdez I-C, Salazar-Henao JE, Fornalé S, López-Vidriero I, Franco-Zorrilla J-M, Grotewold E, Gray J, Solano R, Schmidt W, Pagés M *et al.* 2015. A MYB/ZML complex regulates wound-induced lignin genes in maize. *Plant Cell* 27: 3245–3259.
- Vermaas JV, Dixon RA, Chen F, Mansfield SD, Boerjan W, Ralph J, Crowley MF, Beckham GT. 2019. Passive membrane transport of lignin-related compounds. *Proceedings of the National Academy of Sciences, USA* 116: 23117–23123.
- Wang H, Avci U, Nakashima J, Hahn MG, Chen F, Dixon RA. 2010. Mutation of WRKY transcription factors initiates pith secondary wall formation and increases stem biomass in dicotyledonous plants. *Proceedings of the National Academy of Sciences, USA* 107: 22338–22343.
- Wang S, Yang H, Mei J, Liu X, Wen Z, Zhang L, Xu Z, Zhang B, Zhou Y. 2019. A rice homeobox protein KNAT7 integrates the pathways regulating cell expansion and wall stiffness. *Plant Physiology* 181: 669–682.
- Wang Y, Bouchabke-Coussa O, Lebris P, Antelme S, Soulhat C, Gineau E, Dalmis M, Bendahmane A, Morin H, Mouille G *et al.* 2015. LACCASE5s is required for lignification of the *Brachypodium distachyon* culm. *Plant Physiology* 168: 192.
- Wang Y, Chantreau M, Sibout R, Hawkins S. 2013. Plant cell wall lignification and monolignol metabolism. *Frontiers in Plant Science* 4: 220.
- Whitehead C, Ostos Garrido FJ, Reymond M, Simister R, Distelfeld A, Atienza SG, Piston F, Gomez LD, McQueen-Mason SJ. 2018. A glycosyl transferase family 43 protein involved in xylan biosynthesis is associated with straw digestibility in *Brachypodium distachyon*. *New Phytologist* 218: 974–985.
- Wilkerson CG, Mansfield SD, Lu F, Withers S, Park J-Y, Karlen SD, Gonzales-Vigil E, Padmakshan D, Unda F, Rencoret J *et al.* 2014. Monolignol ferulate transferase introduces chemically labile linkages into the lignin backbone. *Science* 344: 90–93.
- Wilson SM, Burton RA, Doblin MS, Stone BA, Newbigin EJ. 2006. Temporal and spatial appearance of wall polysaccharides during cellularization of barley (*Hordeum vulgare*) endosperm. *Planta* 224: 655.
- Wilson SM, Ho YY, Lampugnani ER, Van de Meene AML, Bain MP, Bacic A, Doblin MS. 2015. Determining the subcellular location of synthesis and assembly of the cell wall polysaccharide (1,3; 1,4)- $\beta$ -D-glucan in grasses. *Plant Cell* 27: 754–771.
- Withers S, Lu F, Kim H, Zhu Y, Ralph J, Wilkerson CG. 2012. Identification of grass-specific enzyme that acylates monolignols with *p*-coumarate. *The Journal of Biological Chemistry* 287: 8347–8355.
- Yang L, Zhao X, Yang F, Fan D, Jiang Y, Luo K. 2016. PtrWRKY19, a novel WRKY transcription factor, contributes to the regulation of pith secondary wall formation in *Populus trichocarpa*. *Scientific Reports* 6: 18643.
- Ye Y, Liu B, Zhao M, Wu K, Cheng W, Chen X, Liu Q, Liu Z, Fu X, Wu Y. 2015. CEF1/OsMYB103L is involved in GA-mediated regulation of secondary wall biosynthesis in rice. *Plant Molecular Biology* 89: 385–401.
- Zhang B, Zhang L, Li F, Zhang D, Liu X, Wang H, Xu Z, Chu C, Zhou Y. 2017. Control of secondary cell wall patterning involves xylan deacetylation by a GDSL esterase. *Nature Plants* 3: 17017.
- Zhang D, Xu Z, Cao S, Chen K, Li S, Liu X, Gao C, Zhang B, Zhou Y. 2018. An uncanonical CCCH-tandem zinc-finger protein represses secondary wall synthesis and controls mechanical strength in rice. *Molecular Plant* 11: 163–174.
- Zhang J, Xie M, Tuskan GA, Muchero W, Chen J-G. 2018. Recent advances in the transcriptional regulation of secondary cell wall biosynthesis in the woody plants. *Frontiers in Plant Science* 9: 1535.
- Zhao K, Bartley L. 2014. Comparative genomic analysis of the R2R3 MYB secondary cell wall regulators of *Arabidopsis*, poplar, rice, maize and switchgrass. *BMC Plant Biology* 14: 135.
- Zhao K, Lin F, Romero-Gamboa SP, Goh H-J, Saha P, An G, Jung K-H, Hazen SP, Bartley LE. 2019. Rice genome-scale network integration reveals transcriptional regulators of grass cell wall synthesis. *Frontiers in Plant Science* 10: 1275.
- Zhong R, Cui D, Dasher RL, Ye Z-H. 2018a. Biochemical characterization of rice xylan *O*-acetyltransferases. *Planta* 247: 1489–1498.
- Zhong R, Cui D, Phillips DR, Ye Z-H. 2018b. A novel rice xylosyltransferase catalyzes the addition of 2-*O*-xylosyl side chains onto the xylan backbone. *Plant & Cell Physiology* 59: 554–565.
- Zhong R, Cui D, Ye Z-H. 2017. Regiospecific acetylation of xylan is mediated by a group of DUF231-containing *O*-acetyltransferases. *Plant & Cell Physiology* 58: 2126–2138.
- Zhong R, Demura T, Ye ZH. 2006. SND1, a NAC domain transcription factor, is a key regulator of secondary wall synthesis in fibers of *Arabidopsis*. *Plant Cell* 18: 3158–3170.
- Zhong R, Lee C, McCarthy RL, Reeves CK, Jones EG, Ye Z-H. 2011. Transcriptional activation of secondary wall biosynthesis by rice and maize NAC and MYB transcription factors. *Plant & Cell Physiology* 52: 1856–1871.
- Zhong R, Lee C, Zhou J, McCarthy RL, Ye Z-H. 2008. A battery of transcription factors involved in the regulation of secondary cell wall biosynthesis in *Arabidopsis*. *Plant Cell* 20: 2763–2782.
- Zhong R, Richardson EA, Ye ZH. 2007. The MYB46 transcription factor is a direct target of SND1 and regulates secondary wall biosynthesis in *Arabidopsis*. *Plant Cell* 19: 2776–2792.
- Zhong R, Ye Z-H. 2007. Regulation of cell wall biosynthesis. *Current Opinion in Plant Biology* 10: 564–572.
- Zhong R, Ye Z-H. 2012. MYB46 and MYB83 bind to the SMRE sites and directly activate a suite of transcription factors and secondary wall biosynthetic genes. *Plant & Cell Physiology* 53: 368–380.
- Zhong R, Yuan Y, Spiekerman JJ, Guley JT, Egboisiuba JC, Ye Z-H. 2015. Functional characterization of NAC and MYB transcription factors involved in regulation of biomass production in switchgrass (*Panicum virgatum*). *PLoS ONE* 10: e0134611.

## Supporting Information

Additional Supporting Information may be found online in the Supporting Information section at the end of the article.

**Table S1** List of *B. distachyon* genes related to secondary cell wall synthesis.

Please note: Wiley Blackwell are not responsible for the content or functionality of any Supporting Information supplied by the authors. Any queries (other than missing material) should be directed to the *New Phytologist* Central Office.